

L^p bounds for a combinatorial central limit theorem with involutions

Subhankar Ghosh*

University of Southern California

April 22, 2010

Abstract

Let $E = ((e_{ij}))_{n \times n}$ be a fixed array of real numbers such that $e_{ij} = e_{ji}$, $e_{ii} = 0$ for $1 \leq i, j \leq n$. Let the symmetric group be denoted by S_n and the collection of involutions with no fixed points by Π_n , that is, $\Pi_n = \{\pi \in S_n : \pi^2 = id, \pi(i) \neq i \forall i\}$. For π uniformly chosen from Π_n , let $Y_E = \sum_{i=1}^n e_{i\pi(i)}$ and $W = (Y_E - \mu_E)/\sigma_E$ where $\mu_E = E(Y_E)$ and $\sigma_E^2 = \text{Var}(Y_E)$. Denoting by F_W and Φ the distribution functions of W and a $\mathcal{N}(0, 1)$ variate respectively, we bound $\|F_W - \Phi\|_p$ for $1 \leq p \leq \infty$ using Stein's method and zero bias transformations. The resulting bound obtained is the product of a third moment type quantity multiplied by an explicit constant, and in particular for $p = \infty$ is of the same form as the one obtained by Bolthausen for Hoeffding's combinatorial central limit theorem when π is chosen uniformly from S_n . The approximation developed here for involutions has applications in testing whether there is a significant degree of similarity in certain matched pairs experiments.

1 Introduction

Let $E = ((e_{ij}))$ be an $n \times n$ array of real numbers. The study of combinatorial central limit theorems, that is, central limit theorems for random variables of the form

$$Y_E = \sum_{i=1}^n e_{i\pi(i)} \quad (1)$$

usually focuses on the case where π is a permutation chosen uniformly from the set of all permutations of n elements. Approximating the distribution of Y_E by the normal when π is chosen uniformly began with the work of Wald and Wolfowitz [14], who, motivated by approximating null distributions for permutation test statistics, proved the central limit theorem as $n \rightarrow \infty$ for the case where the factorization $e_{ij} = b_i c_j$ holds. In this special case, when b_i are numerical characteristics of some population, and k of the c_j 's are 1, and the remaining $n - k$ are zero, Y_E has the distribution of a simple random sample from the population. General arrays were handled in the work of Hoeffding [8], and in Motoo obtained Lindeberg-type conditions which are sufficient for the normal limit in [10].

A number of later authors refined the limiting results and obtained information on the rate of convergence and bounds on the error in the normal approximation, typically in the supremum, or L^∞ , norm. Ho and Chen [7] and von Bahr [13] derived L^∞ bounds when the matrix E is random, the former using a concentration inequality approach and Stein's method, but which yield the correct rate $O(n^{-1/2})$ only when certain boundedness conditions on $\sup_{i,j} |e_{i,j}|$ are satisfied. Bolthausen [1] obtained a bound of the right order in terms of third-moment type quantities on E , but with an unspecified constant. Goldstein [4] employing

*Department of Mathematics, University of Southern California, Los Angeles, CA 90089, USA, subhankg@usc.edu

the zero bias version of Stein's method obtained bounds of the correct order with an explicit constant, but in terms of $\sup_{i,j} |e_{i,j}|$. Overall then, the work of Bolthausen [1], proceeding inductively, is the only one which yields a bound in terms of third moment quantities on E without the need for conditions on $\sup_{i,j} |e_{i,j}|$. Though an explicit constant was not obtained in [1], as we will see here for a related case, this inductive method can yield explicit constants when the calculations are tracked carefully.

The case where π is chosen uniformly from the symmetric group is not the only case of practical interest. In [11] a permutation test is considered for a certain matched pair experiment, which is designed to answer the question of whether there is an unusually high degree of similarity in a distinguished pairing, when there is some unknown background, or baseline, similarity between all pairs. In such a case one considers Y_E as in (1) when π is chosen uniformly from Π_n , the class of involutions without fixed points,

$$\Pi_n = \{\pi \in S_n : \pi^2 = \text{id}, \forall i : \pi(i) \neq i\}, \quad (2)$$

where S_n is the symmetric group of some n elements and id the identity permutation. Since the distribution of Y_E is complicated, L^∞ bounds for the error in normal approximation enables one to test the significance of a matched pair. The interested reader can look into [6] for a discussion on similar applications.

A bound to the normal for this case was provided in the L^∞ norm by [6], with explicit constants, but under a boundedness assumption. In this paper, we follow the method of [1] to refine the results of [6] and provide a bound to the normal for the involution case, with an explicit constant, in terms of third moment type quantities on the matrix E , and for all L^p for $1 \leq p \leq \infty$. In particular, with K_p given explicitly by (14), we show that the L^p norm of the difference between W , the variable Y_E standardized, and the normal satisfies

$$\|F_W - \Phi\|_p \leq K_p \frac{\beta_E}{n}$$

for all $n \geq 9$, where β_E is given in (11). Although the constant K_p is quite large in magnitude, it is the first of its kind in the literature. Also we improve upon Goldstein and Rinott's result by using the idea of zero bias transformations introduced in [5] and prove CLT of order $O(n^{-1/2})$ under much milder conditions. It should be noted that the method applied here can be adopted to give L^p estimates in Hoeffding's combinatorial CLT as well, and will yield a bound of the same form as the one obtained in [1].

Other work where π is not uniform includes that of [9], where the permutations are uniform over those having one long cycle, and [4], where L^∞ bounds are derived under a boundedness condition for the case of a permutation distribution constant on conjugacy classes having no fixed points, which generalizes both the involution and long cycle cases.

The paper is organized as follows. In Section 2, we introduce some notation and state our main result. In Section 3 the basic idea of the zero bias transformation is reviewed, and an outline is provided that illustrates how to obtain zero bias couplings in some cases of interest. In Section 4, L^1 bounds are obtained. Lastly, in Section 5 we use the calculations in Section 4 along with the recursive argument of [1] to obtain L^∞ bounds. From the L^1 and L^∞ bounds, the following simple inequality allows for the computation of L^p bounds for all intermediate $p \in (1, \infty)$,

$$\|f\|_p^p \leq \|f\|_\infty^{p-1} \|f\|_1. \quad (3)$$

2 Notation and statement of main result

For n an even positive integer, let π be a permutation chosen uniformly from Π_n in (2), the set of involutions with no fixed points. Since for $\pi \in \Pi_n$ the terms $e_{i\pi(i)}$ and $e_{\pi(i)i}$ always appear together in (1), and e_{ii} never appears, we may assume without loss of generality that

$$e_{ij} = e_{ji} \quad \text{and that} \quad e_{ii} = 0 \quad \text{for all } i, j = 1, 2, \dots, n. \quad (4)$$

For an array $E = ((e_{ij}))_{1 \leq i, j \leq n}$ satisfying the conditions in (4), define

$$e_{i+} = \sum_{j=1}^n e_{ij}, \quad e_{+j} = \sum_{i=1}^n e_{ij} \quad \text{and} \quad e_{++} = \sum_{i,j=1}^n e_{ij}.$$

Then, as shown in [6],

$$\begin{aligned}\mu_E &= EY_E = \frac{e_{++}}{n-1} \\ \sigma_E^2 &= \frac{2}{(n-1)(n-3)} \left((n-2) \sum_{1 \leq i, j \leq n} e_{ij}^2 + \frac{1}{n-1} e_{++}^2 - 2 \sum_{i=1}^n e_{i+}^2 \right).\end{aligned}\quad (5)$$

Again following [6], letting

$$\hat{e}_{ij} = \begin{cases} e_{ij} - \frac{e_{i+}}{n-2} - \frac{e_{+j}}{n-2} + \frac{e_{++}}{(n-1)(n-2)} & i \neq j \\ 0 & i = j, \end{cases}\quad (6)$$

we have

$$\hat{e}_{+i} = \hat{e}_{j+} = \hat{e}_{++} = 0 \quad \text{for all } i, j = 1, \dots, n. \quad (7)$$

and that

$$Y_{\hat{E}} = Y_E - \mu_E$$

where $\hat{E}$ is obtained from E by (6).

We consider bounds to the normal $\mathcal{N}(0, 1)$ for the standardized variable

$$W = \frac{Y_E - \mu_E}{\sigma_E}. \quad (8)$$

Since Y_E and $Y_{\hat{E}}$ differ by a constant, (5) and (7) yield

$$\sigma_E^2 = \sigma_{\hat{E}}^2 = \frac{2(n-2)}{(n-1)(n-3)} \sum_{1 \leq i, j \leq n} \hat{e}_{ij}^2. \quad (9)$$

In particular, the mean zero, variance 1 random variable W in (8) can be written as

$$W = \sum_{i=1}^n d_{i\pi(i)} \quad \text{where } d_{ij} = \hat{e}_{ij}/\sigma_E, \quad (10)$$

and moreover, the array $\hat{E}$ inherits properties (4) and (7) from E , as then does D from $\hat{E}$. For any array $E = ((e_{ij}))_{n \times n}$, let

$$\beta_E = \sum_{i \neq j} \frac{|\hat{e}_{ij}|^3}{\sigma_E^3} \quad \text{with } \hat{e}_{ij} \text{ as in (6) and } \sigma_E^2 \text{ as in (5)}. \quad (11)$$

In order to guarantee that W is a well defined random variable, that is, that $\sigma_E^2 > 0$, we henceforth impose the following condition without further mention.

Condition 2.1. *For some $i \neq j$, the value $\hat{e}_{ij} \neq 0$, or, equivalently,*

$$e_{ij} - \frac{e_{i+}}{n-2} - \frac{e_{+j}}{n-2} + \frac{e_{++}}{(n-1)(n-2)} \neq 0 \quad \text{for some } i \neq j.$$

Since d_{ij} and $\hat{e}_{ij}$ are linearly related, Condition 2.1 is equivalent to the condition that $d_{ij} \neq 0$ for some $i \neq j$.

We provide bounds on the accuracy of the normal approximation of W using the zero bias transformation, introduced in [5]. For any random variable W with mean zero and variance σ^2 , there exists a unique distribution for a random variable W^* with the property that for any differentiable function f

$$E[Wf(W)] = \sigma^2 E f'(W^*). \quad (12)$$

The distribution of the random variate W^* is called the zero bias transform of the distribution of W . From Stein's original lemma [12], W is $\mathcal{N}(0, \sigma^2)$ if and only if W^* is $\mathcal{N}(0, \sigma^2)$, that is, the normal distribution is the unique fixed point of the zero bias transformation. This gives rise to the intuition that if W and W^* are close, in some appropriate sense, then the distribution of W should be close to the normal. That this intuition is indeed true can be seen, for instance, in the following result from [3]: if W and W^* are on a joint space, with W^* having the W zero bias distribution, then

$$\|F_W - \Phi\|_1 \leq 2E|W^* - W|. \quad (13)$$

In (13), F_W and Φ denote the distribution functions of W and that of a standard normal variate respectively, and $\|\cdot\|_p$ denotes the L^p norm. We call a construction of W and W^* on a joint space a zero bias coupling of W to W^* .

The following is our main result which we prove using zero bias coupling.

Theorem 2.1. *Let $E = ((e_{ij}))_{n \times n}$ be an array satisfying $e_{ij} = e_{ji}$, $e_{ii} = 0 \forall i, j$, and let π be an involution chosen uniformly from Π_n . If*

$$Y_E = \sum_{i=1}^n e_{i\pi(i)},$$

and $W = (Y_E - \mu_E)/\sigma_E$, then for $n \geq 9$ and $p \in [1, \infty]$, with β_E as in (11), we have

$$\|F_W - \Phi\|_p \leq K_p \frac{\beta_E}{n}.$$

Here F_W denotes the distribution function of W , Φ is the distribution function of a standard normal variate and

$$K_p = (379)^{1/p} (61, 702, 446)^{1-1/p}. \quad (14)$$

As W in the theorem is given by (10) with

$$d_{ij} = d_{ji}, d_{ii} = 0, d_{i+} = 0 \quad \text{and} \quad \sigma_D^2 = 1; \beta_E = \beta_D \quad (15)$$

we assume in what follows that all subsequent occurrences of $((d_{ij}))$ satisfy these conditions and instead of working with E work with the centered and scaled array D only.

In the next section, we review construction of zero bias couplings in certain cases of interest including the present problem.

3 Zero bias transformation

We prove Theorem 2.1 by constructing a zero bias coupling using a Stein pair, that is, a pair of random variables (W, W') which are exchangeable and satisfy

$$E(W - W'|W) = \lambda W \quad \text{for some } \lambda \in (0, 1); \quad (16)$$

see [2] for more on Stein pairs.

As shown in [5], for any mean zero, variance σ^2 random variable W , there exists a distribution for a random variable W^* satisfying (12). Nevertheless, constructing useful couplings of W and W^* for particular examples may be difficult. In some cases, however, as in ours, the following proposition from [3] may be applied.

Proposition 3.1. *Let W, W' be an exchangeable pair with $\text{Var}(W) = \sigma^2 \in (0, \infty)$ and distribution $F(w, w')$ satisfying the linearity condition (16). Then*

$$E(W - W')^2 = 2\lambda\sigma^2, \quad (17)$$

and when $(W^\dagger, W^\ddagger)$ has the joint distribution

$$dF^\dagger(w, w') = \frac{(w - w')^2}{E(W - W')^2} dF(w, w') \quad (18)$$

and $U \sim \mathcal{U}[0, 1]$ is independent of $W^\dagger, W^\ddagger$, the variable

$$W^* = UW^\dagger + (1 - U)W^\ddagger$$

has the W -zero biased distribution.

In particular, construction of zero bias couplings are typically possible when Stein pairs exist. We review the construction of $(W^\dagger, W^\ddagger)$ from (W, W') as outlined in [3]. Suppose we have a Stein pair (W, W') which is a function of some collection of random variables $\{\Xi_\alpha, \alpha \in \chi\}$, and that for a possibly random index set $\mathbf{I} \subset \chi$, independent of $\{\Xi_\alpha, \alpha \in \chi\}$, the difference $W - W'$ depends only on $\mathbf{I}$ and on $\{\Xi_\alpha, \alpha \in \chi_{\mathbf{I}}\}$, where $\chi_{\mathbf{I}} \subset \chi$ depends on $\mathbf{I}$. That is, for some function $b(\mathbf{i}, \Xi_\alpha, \alpha \in \chi_{\mathbf{i}})$ defined on $\mathbf{i} \subset \chi$, and $\mathbf{I}$ a random index set,

$$W - W' = b(\mathbf{I}, \Xi_\alpha, \alpha \in \chi_{\mathbf{I}}). \quad (19)$$

We now show how, under this framework, the pair (W, W') can be constructed; the pair $(W^\dagger, W^\ddagger)$ will then be constructed in a similar fashion. First generate $\mathbf{I}$, then independently generate $\{\Xi_\alpha, \alpha \in \chi_{\mathbf{I}}\}$ and finally $\{\Xi_\alpha, \alpha \in \chi_{\mathbf{I}}^c\}$ conditioned on $\{\Xi_\alpha, \alpha \in \chi_{\mathbf{I}}\}$. That is, first generate the indices $\mathbf{I}$ on which the difference $W - W'$ depends, then the underlying variables $\Xi_\alpha, \alpha \in \chi_{\mathbf{I}}$ which make up that difference, and lastly the remaining variables. This construction corresponds to the following factorization of the joint distribution of $\mathbf{I}$ and $\{\Xi_\alpha, \alpha \in \chi\}$ as the product

$$dF(\mathbf{i}, \xi_\alpha, \alpha \in \chi) = P(\mathbf{I} = \mathbf{i}) dF_{\mathbf{i}}(\xi_\alpha, \alpha \in \chi_{\mathbf{i}}) dF_{\mathbf{i}^c|\mathbf{i}}(\xi_\alpha, \alpha \notin \chi_{\mathbf{i}} | \xi_\alpha, \alpha \in \chi_{\mathbf{i}}). \quad (20)$$

For $dF^\dagger$ we consider the joint distribution of $\mathbf{I}$ and $\{\Xi_\alpha, \alpha \in \chi\}$, biased by the squared difference $(w - w')^2$, is

$$dF^\dagger(\mathbf{i}, \xi_\alpha, \alpha \in \chi) = \frac{(w - w')^2}{E(W - W')^2} dF(\mathbf{i}, \xi_\alpha, \alpha \in \chi). \quad (21)$$

From (17), (19) and the independence of $\mathbf{I}$ and $\{\Xi_\alpha, \alpha \in \chi\}$ we obtain

$$\sum_{\mathbf{i} \subset \chi} P(\mathbf{I} = \mathbf{i}) E b^2(\mathbf{i}, \Xi_\alpha, \alpha \in \chi_{\mathbf{i}}) = 2\lambda\sigma^2. \quad (22)$$

Hence we can define a probability distribution for an index set $\mathbf{I}^\dagger$ by

$$P(\mathbf{I}^\dagger = \mathbf{i}) = \frac{r_{\mathbf{i}}}{2\lambda\sigma^2} \quad \text{where} \quad r_{\mathbf{i}} = P(\mathbf{I} = \mathbf{i}) E b^2(\mathbf{i}, \Xi_\alpha, \alpha \in \chi_{\mathbf{i}}). \quad (23)$$

From (19), (21) and (23), we obtain

$$\begin{aligned} dF^\dagger(\mathbf{i}, \xi_\alpha, \alpha \in \chi) &= \frac{b^2(\mathbf{i}, \xi_\alpha, \alpha \in \chi_{\mathbf{i}})}{2\lambda\sigma^2} P(\mathbf{I} = \mathbf{i}) dF_{\mathbf{i}}(\xi_\alpha, \alpha \in \chi_{\mathbf{i}}) dF_{\mathbf{i}^c|\mathbf{i}}(\xi_\alpha, \alpha \notin \chi_{\mathbf{i}} | \xi_\alpha, \alpha \in \chi_{\mathbf{i}}) \\ &= \frac{r_{\mathbf{i}}}{2\lambda\sigma^2} \frac{b^2(\mathbf{i}, \xi_\alpha, \alpha \in \chi_{\mathbf{i}})}{E b^2(\mathbf{i}, \Xi_\alpha, \alpha \in \chi_{\mathbf{i}})} dF_{\mathbf{i}}(\xi_\alpha, \alpha \in \chi_{\mathbf{i}}) dF_{\mathbf{i}^c|\mathbf{i}}(\xi_\alpha, \alpha \notin \chi_{\mathbf{i}} | \xi_\alpha, \alpha \in \chi_{\mathbf{i}}) \\ &= P(\mathbf{I}^\dagger = \mathbf{i}) dF_{\mathbf{i}}^\dagger(\xi_\alpha, \alpha \in \chi_{\mathbf{i}}) dF_{\mathbf{i}^c|\mathbf{i}}(\xi_\alpha, \alpha \notin \chi_{\mathbf{i}} | \xi_\alpha, \alpha \in \chi_{\mathbf{i}}) \end{aligned} \quad (24)$$

where

$$dF_{\mathbf{i}}^\dagger(\xi_\alpha, \alpha \in \chi_{\mathbf{i}}) = \frac{b^2(\mathbf{i}, \xi_\alpha, \alpha \in \chi_{\mathbf{i}})}{E b^2(\mathbf{i}, \Xi_\alpha, \alpha \in \chi_{\mathbf{i}})} dF_{\mathbf{i}}(\xi_\alpha, \alpha \in \chi_{\mathbf{i}}). \quad (25)$$

Note that (24) gives a representation of the distribution $F^\dagger$ which is of the same form as (20). The parallel forms of F and $F^\dagger$ allow us to generate variables having distribution $F^\dagger$ parallel to that for distribution F ; first generate the random index $\mathbf{I}^\dagger$, then $\{\Xi_\alpha^\dagger, \alpha \in \chi_{\mathbf{I}^\dagger}\}$ according to $dF_{\mathbf{I}^\dagger}^\dagger$, and lastly the remaining variables according to $dF_{\mathbf{I}^c|\mathbf{I}^\dagger}(\xi_\alpha, \alpha \notin \chi_{\mathbf{I}^\dagger} | \xi_\alpha, \alpha \in \chi_{\mathbf{I}^\dagger})$.

That the last step is the same in the construction of pairs of variables having the F and $F^\dagger$ distribution allows an opportunity for a coupling between (W, W') and $(W^\dagger, W^{\ddagger})$ to be achieved; from Proposition 3.1, this suffices to couple W and W^* . One coupling may be accomplished using the following outline, as was done in [3]. First generate $\mathbf{I}$ and $\{\Xi_\alpha, \alpha \in \chi\}$, yielding the pair W, W' . Next generate $\mathbf{I}^\dagger$ and then the variables $\{\Xi_\alpha^\dagger : \alpha \in \chi_{\mathbf{I}^\dagger}\}$ following $dF_{\mathbf{I}^\dagger}^\dagger$ if the realization of $\mathbf{I}^\dagger$ is $\mathbf{i}$. Lastly, when constructing the remaining variables that is $\{\Xi_\alpha^\dagger : \alpha \notin \chi_{\mathbf{i}}\}$ which make up $W^\dagger, W^{\ddagger}$, use as much of the previously generated variables $\{\Xi_\alpha, \alpha \notin \chi_{\mathbf{i}}\}$ as possible so that the pairs (W, W') and $(W^\dagger, W^{\ddagger})$ will be close.

We review the construction of (W, W') in [6] for the case at hand, and then show how it agrees with the outline above. For distinct $i, j \in \{1, \dots, n\}$ let $\tau_{i,j}$ be the permutation which transposes the elements i and j , that is, $\tau_{i,j}(i) = j, \tau_{i,j}(j) = i$ and $\tau_{i,j}(k) = k$ for all $k \notin \{i, j\}$. Furthermore, given $\pi \in \Pi_n$, let

$$\alpha_{i,j}^\pi = \tau_{i,\pi(j)}\tau_{j,\pi(i)}.$$

Note that for any given $\pi \in \Pi_n$, the permutation $\pi\alpha_{i,j}^\pi$ will again belong to Π_n . In particular, whereas π has the cycle(s) $(i, \pi(i))$ and $(j, \pi(j))$, the permutation $\pi\alpha_{i,j}^\pi$ has cycle(s) (i, j) and $(\pi(i), \pi(j))$, and all other cycles in common with π . Now, with π chosen uniformly from Π_n and W given by (10), we construct an exchangeable pair (W, W') , as in [6], as follows. Choose two distinct indices $\mathbf{I} = (I, J)$ uniformly from $\{1, 2, \dots, n\}$, that is, having distribution

$$P(\mathbf{I} = \mathbf{i}) = P(I = i, J = j) = \frac{1}{n(n-1)} \mathbf{1}(i \neq j) \quad (26)$$

and let

$$\pi' = \pi\alpha_{I,J}^\pi. \quad (27)$$

Since (I, J) is chosen uniformly over all distinct pairs, π and π' are exchangeable, and hence, letting $W' = \sum d_{i\pi'(i)}$, so are (W, W') . Moreover, as π' has the cycle(s) (I, J) and $(\pi(I), \pi(J))$, and shares all other cycles with π , we have

$$W - W' = 2(d_{I\pi(I)} + d_{J\pi(J)} - (d_{IJ} + d_{\pi(I)\pi(J)})). \quad (28)$$

Using (28) it is shown in [6] that

$$E(W - W'|W) = \frac{4}{n}W, \quad (29)$$

that is, (16) is satisfied with $\lambda = 4/n$.

To put this construction in the framework above, so to be able to apply the decomposition (24) for the construction of a zero bias coupling, let $\chi = \{1, 2, \dots, n\}$ and $\Xi_\alpha = \pi(\alpha)$. From (28), we see that the difference $W - W'$ depends on a pair of randomly chosen indices, and their images. Hence, regarding these indices, let $\mathbf{i} = (i, j) \in \chi^2$ and let $\mathbf{I} = (I, J)$ where I and J have joint distribution given in (26), specifying $P(\mathbf{I} = \mathbf{i})$, the first term in (20). Also, in this case $\chi_{\mathbf{i}} = \mathbf{i}$. Next, for given distinct i, j and $\pi \in \Pi_n$, we have $\pi(i) \neq \pi(j), \pi(i) \neq i$ and $\pi(j) \neq j$. As π is chosen uniformly from Π_n , the distribution of the images $k = \xi_i$ and $l = \xi_j$ of distinct i and j under π is given by

$$dF_{\mathbf{i}}(\xi_\alpha, \alpha \in \mathbf{i}) = dF_{i,j}(k, l) \propto \mathbf{1}(k \neq l, k \neq i, l \neq j) \quad \text{for } i \neq j, \quad (30)$$

specifying the second term of (20). The last term in (20) is given by

$$dF_{\mathbf{i}^c|\mathbf{i}}(\xi_\alpha, \alpha \notin \mathbf{i} | \xi_\alpha, \alpha \in \mathbf{i}) = \frac{P(\pi(i) = k, \pi(j) = l, \pi(\alpha) = \xi_\alpha, \alpha \notin \{i, j, k, l\})}{P(\pi(i) = k, \pi(j) = l)}, \quad (31)$$

when $i \neq j$ and $k \neq l$. The equality in (31) follows from the fact that π is an involution, implying $\xi_k = \pi(k) = \pi(\pi(i)) = i$ and similarly for $\xi_l = j$, and thus

$$\begin{aligned} & \{\pi(i) = k, \pi(j) = l, \pi(\alpha) = \xi_\alpha, \alpha \notin \{i, j\}\} \\ &= \{\pi(i) = k, \pi(j) = l, \pi(k) = i, \pi(l) = j, \pi(\alpha) = \xi_\alpha, \alpha \notin \{i, j, k, l\}\} \\ &= \{\pi(i) = k, \pi(j) = l, \pi(\alpha) = \xi_\alpha, \alpha \notin \{i, j, k, l\}\}. \end{aligned}$$

The conditional distribution in (31) can be simplified further by considering the two cases $k = j$ or equivalently $l = i$ and $k \neq j$ or equivalently $|\{i, j, k, l\}| = 4$ separately. If $k = j$ then we have $l = i$ and thus we obtain

$$\begin{aligned} dF_{\mathbf{i}^c|\mathbf{i}}(\xi_\alpha, \alpha \notin \mathbf{i} | \xi_i = j, \xi_j = i) &= \frac{P(\pi(i) = j, \pi(j) = i, \pi(\alpha) = \xi_\alpha, \alpha \notin \{i, j\})}{P(\pi(i) = j, \pi(j) = i)} \\ &= \frac{P(\pi(i) = j, \pi(\alpha) = \xi_\alpha, \alpha \notin \{i, j\})}{P(\pi(i) = j)} \\ &= \frac{|\Pi_n|^{-1}}{(n-1)^{-1}} = |\Pi_{n-2}|^{-1}. \end{aligned} \quad (32)$$

For (32), we note $P(\pi(i) = j) = 1/(n-1)$ since π is chosen uniformly from Π_n . The last equality in (32) simply indicates that once we fix $\pi(i) = j$, that is the cycle (i, j) in π we have to choose an involution uniformly at random from the rest of the indices that is Π_{n-2} to obtain π in its entirety. This argument yields the recursion which we use in (33) later

$$|\Pi_n| = (n-1)|\Pi_{n-2}|.$$

When we have $k \neq j$ and hence $l \neq i$ or equivalently $|\{i, j, k, l\}| = 4$ in (31), we obtain

$$\begin{aligned} dF_{\mathbf{i}^c|\mathbf{i}}(\xi_\alpha, \alpha \notin \mathbf{i} | \xi_\alpha, \alpha \in \mathbf{i}) &= \frac{P(\pi(i) = k, \pi(j) = l, \pi(\alpha) = \xi_\alpha, \alpha \notin \{i, j, k, l\})}{P(\pi(i) = k, \pi(j) = l)} \\ &= \frac{|\Pi_n|^{-1}}{((n-1)(n-3))^{-1}} = |\Pi_{n-4}|^{-1}. \end{aligned} \quad (33)$$

In (33) we have used the following equality which follows from the fact that π is an involution chosen uniformly at random.

$$P(\pi(i) = k, \pi(j) = l) = P(\pi(i) = k)P(\pi(j) = l | \pi(i) = k) = \frac{1}{(n-3)(n-1)}.$$

From (33) we see that the conditional distribution $dF_{\mathbf{i}^c|\mathbf{i}}$ is uniform over all values of ξ_α for $\alpha \in \chi$ for which $\xi_i = k, \xi_j = l$ and $P(\pi(\alpha) = \xi_\alpha, \alpha \in \chi) > 0$ when $|\{i, j, k, l\}| = 4$. Hence we may construct (W, W') following (20), that is, first choosing $\mathbf{I} = (I, J)$, then the images $\mathbf{K} = (K, L)$ of I and J under π and π' , then the remaining images uniformly over all possible values for which the resulting permutations lie in Π_n .

We may construct $(W^\dagger, W^\ddagger)$ from (24) quite easily now. In view of (19) and (28), for pairs of distinct indices i, j , let

$$b(i, j, \pi_i, \pi_j) = 2(d_{i\pi_i} + d_{j\pi_j} - (d_{ij} + d_{\pi_i\pi_j})), \quad (34)$$

where again we may also let $k = \xi_i$ and $l = \xi_j$. Now, considering the first two factors in (24), using (23), (25), (26) and (30) we obtain,

$$\begin{aligned} P(\mathbf{I}^\dagger = \mathbf{i})dF_{\mathbf{i}^\dagger}(\xi_\alpha, \alpha \in \mathbf{i}) &= P(I^\dagger = i, J^\dagger = j)dF_{i,j}^\dagger(K^\dagger = k, L^\dagger = l) \\ &\propto P(I = i, J = j)[d_{ik} + d_{jl} - (d_{ij} + d_{kl})]^2 \mathbf{1}(k \neq l, k \neq i, l \neq j) \\ &\propto [d_{ik} + d_{jl} - (d_{ij} + d_{kl})]^2 \mathbf{1}(k \neq l, k \neq i, l \neq j, i \neq j). \end{aligned} \quad (35)$$

By Lemma 3.3, below, relation (35) specifies a joint distribution, say $p(i, j, k, l)$, on the pairs $\mathbf{I}^\dagger = (I^\dagger, J^\dagger)$ and their images $(\xi_{I^\dagger}, \xi_{J^\dagger}) = (K^\dagger, L^\dagger) := \mathbf{K}^\dagger$, say. Since when $j = k$, or equivalently $i = l$ the square term vanishes, we may write

$$p(i, j, k, l) = c_n [d_{ik} + d_{jl} - (d_{ij} + d_{kl})]^2 \mathbf{1}(|\{i, j, k, l\}| = 4), \quad (36)$$

where the constant of proportionality c_n is provided in Lemma 3.3. Next, note that since i, j, k, l have to be distinct in the definition (36), the third term in (24), $dF_{\mathbf{i}^c|\mathbf{i}}(\xi_\alpha, \alpha \notin \mathbf{i} | \xi_\alpha, \alpha \in \mathbf{i})$, reduces to uniform distribution over all values of ξ_α for $\alpha \in \chi$ for which $\xi_i = k, \xi_j = l$ and $P(\pi(\alpha) = \xi_\alpha, \alpha \in \chi) > 0$ owing to (33). Once we form $(W^\dagger, W^\ddagger)$ following the square bias distribution (18), it's easy to produce W^* which has the W zero bias distribution using Proposition 3.1. We summarize the conclusions above in the following lemma.

Lemma 3.1. *Let $dF(w, w')$ be the joint distribution of a Stein pair (W, W') where*

$$W = \sum_{i=1}^n d_{i\pi(i)} \quad \text{and} \quad W' = \sum_{i=1}^n d_{i\pi'(i)}$$

with π chosen uniformly from Π_n and π' as in (27) with I, J having distribution (26). Then a pair $(W^\dagger, W^\ddagger)$ with the square bias distribution (18) can be constructed by setting

$$W^\dagger = \sum_{i=1}^n d_{i\pi^\dagger(i)} \quad \text{and} \quad W^\ddagger = \sum_{i=1}^n d_{i\pi^\ddagger(i)}$$

where $\pi^\dagger$ and $\pi^\ddagger = \pi^\dagger \alpha_{I^\dagger, J^\dagger}^{\pi^\dagger}$ are constructed by first sampling $\mathbf{I}^\dagger = (I^\dagger, J^\dagger)$ and the respective images $\mathbf{K}^\dagger = (K^\dagger, L^\dagger)$ under $\pi^\dagger$ according to (36) and then selecting the remaining images of $\pi^\dagger$ uniformly from among the choices for which it lies in Π_n . Furthermore if $U \sim \mathcal{U}[0, 1]$ is independent of $(W^\dagger, W^\ddagger)$, then $W^ = UW^\dagger + (1 - U)W^\ddagger$ has the W zero bias distribution. Hence, if π and $\pi^\dagger$ are constructed on a common space, then so are W and W^* .*

Given the permutations π and π' from which the pair (W, W') is constructed, we would like to form $(W^\dagger, W^\ddagger)$ as close to (W, W') as possible, thus making W close to W^* . Towards this end, we follow the construction noted after (24), using many of the already chosen variables which form π and π' to make the two pairs close. In particular, begin the construction of $(W^\dagger, W^\ddagger)$ by choosing $\mathbf{I}^\dagger = (I^\dagger, J^\dagger)$ and $\mathbf{K}^\dagger = (K^\dagger, L^\dagger)$ with joint distribution (36), independent of π and π' . Let $R_1 = |\{\pi(I^\dagger), \pi(J^\dagger)\} \cap \{K^\dagger, L^\dagger\}|$ and $R_2 = |\{\pi(I^\dagger), \pi(K^\dagger)\} \cap \{J^\dagger, L^\dagger\}|$; clearly $R_1, R_2 \in \{0, 1, 2\}$. Define $\pi^\dagger$ by

$$\pi^\dagger = \begin{cases} \pi \alpha_{J^\dagger, L^\dagger}^{\pi} & \text{if } \pi(I^\dagger) = K^\dagger \text{ and } \pi(J^\dagger) \neq L^\dagger & \text{hence } (R_1, R_2) = (1, 0) \\ \pi \alpha_{I^\dagger, K^\dagger}^{\pi} & \text{if } \pi(I^\dagger) \neq K^\dagger \text{ and } \pi(J^\dagger) = L^\dagger & \text{hence } (R_1, R_2) = (1, 0) \\ \pi \alpha_{J^\dagger, K^\dagger}^{\pi} \tau_{I^\dagger, J^\dagger} \tau_{K^\dagger, L^\dagger} & \text{if } \pi(I^\dagger) = L^\dagger \text{ and } \pi(J^\dagger) \neq K^\dagger & \text{hence } (R_1, R_2) = (1, 1) \\ \pi \alpha_{I^\dagger, L^\dagger}^{\pi} \tau_{I^\dagger, J^\dagger} \tau_{K^\dagger, L^\dagger} & \text{if } \pi(I^\dagger) \neq L^\dagger \text{ and } \pi(J^\dagger) = K^\dagger & \text{hence } (R_1, R_2) = (1, 1) \\ \pi \alpha_{K^\dagger, L^\dagger}^{\pi} \tau_{I^\dagger, L^\dagger} \tau_{J^\dagger, K^\dagger} & \text{if } \pi(I^\dagger) = J^\dagger \text{ and } \pi(K^\dagger) \neq L^\dagger & \text{hence } (R_1, R_2) = (0, 1) \\ \pi \alpha_{I^\dagger, J^\dagger}^{\pi} \tau_{I^\dagger, L^\dagger} \tau_{J^\dagger, K^\dagger} & \text{if } \pi(I^\dagger) \neq J^\dagger \text{ and } \pi(K^\dagger) = L^\dagger & \text{hence } (R_1, R_2) = (0, 1) \\ \pi & \text{if } \pi(I^\dagger) = K^\dagger \text{ and } \pi(J^\dagger) = L^\dagger & \text{hence } (R_1, R_2) = (2, 0) \\ \pi \tau_{I^\dagger, L^\dagger} \tau_{J^\dagger, K^\dagger} & \text{if } \pi(I^\dagger) = J^\dagger \text{ and } \pi(K^\dagger) = L^\dagger & \text{hence } (R_1, R_2) = (0, 2) \\ \pi \tau_{I^\dagger, J^\dagger} \tau_{K^\dagger, L^\dagger} & \text{if } \pi(I^\dagger) = L^\dagger \text{ and } \pi(J^\dagger) = K^\dagger & \text{hence } (R_1, R_2) = (2, 2) \\ \pi \alpha_{I^\dagger, K^\dagger}^{\pi} \alpha_{J^\dagger, L^\dagger}^{\pi} & \text{when } R_1 = R_2 = 0. & \end{cases} \quad (37)$$

The partition in display (37) is based on the possible values of (R_1, R_2) ; it does not include the cases where $(R_1, R_2) = (2, 1)$ or $(R_1, R_2) = (1, 2)$ because these two events are impossible. If $R_1 = 2$ and $\pi(I^\dagger) = K^\dagger, \pi(J^\dagger) = L^\dagger$ then $R_2 = 0$ while if $\pi(I^\dagger) = L^\dagger, \pi(J^\dagger) = K^\dagger$ then $R_2 = 2$. Similarly one can rule out $(R_1, R_2) = (1, 2)$. Similar arguments show us that the cases described above are indeed exhaustive.

Clearly any two cases in (37) with differing values of (R_1, R_2) tuple are exclusive. Also, any two cases with the same tuple value, e.g. cases one and two, are also exclusive. For example, in case one we have $\pi(I^\dagger) = K^\dagger$ whereas in case two we have $\pi(I^\dagger) \neq K^\dagger$ making these two cases disjoint. In summary $\pi^\dagger$ is well defined, and this construction specifies the pairs (π, π') and $(\pi^\dagger, \pi^\dagger)$ on the same space. The following lemma shows that the $\pi^\dagger$ so obtained is an involution.

Lemma 3.2. *For $\pi \in \Pi_n$, the permutation $\pi^\dagger$ defined in (37) belongs to Π_n and has the cycles $(I^\dagger, K^\dagger)$ and $(J^\dagger, L^\dagger)$. Moreover, with $\pi^\dagger$ as in Lemma 3.1, the permutations $\pi, \pi^\dagger, \pi^\dagger$ are involutions when restricted to the set $\mathcal{I} = \{I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger), \pi(K^\dagger), \pi(L^\dagger)\}$, and agree on the complement $\mathcal{I}^c$.*

Proof. If we prove $\pi^\dagger$ has the cycles $(I^\dagger, K^\dagger)$ and $(J^\dagger, L^\dagger)$, then from (37) and $\pi^\dagger = \pi \alpha_{I^\dagger, J^\dagger}^{\pi^\dagger}$ we have that $\pi, \pi^\dagger, \pi^\dagger$ all agree on the complement of $\mathcal{I}$, which proves the last claim in the lemma. Note that π maps $\mathcal{I}$ onto itself and is an involution when restricted to $\mathcal{I}$. Therefore π is an involution when restricted to $\mathcal{I}^c$, and hence so are $\pi^\dagger$ and $\pi^\dagger$. So, we only need to prove $\pi^\dagger$ has the cycles as claimed.

Since $\pi^\dagger = \pi \alpha_{I^\dagger, J^\dagger}^{\pi^\dagger}$, it suffices now to show that $\pi^\dagger$ is an involution on $\mathcal{I}$ and has cycles $(I^\dagger, K^\dagger)$ and $(J^\dagger, L^\dagger)$, which can be achieved by examining the cases in (37) one by one. For instance, in case one, where $\pi(I^\dagger) = K^\dagger$ and $\pi(J^\dagger) \neq L^\dagger$ we have $\mathcal{I} = \{I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(J^\dagger), \pi(L^\dagger)\}$ and

$$\begin{aligned} \pi^\dagger(I^\dagger) &= \pi \alpha_{J^\dagger, L^\dagger}^{\pi^\dagger}(I^\dagger) = \pi(I^\dagger) = K^\dagger \\ \pi^\dagger(J^\dagger) &= \pi \alpha_{J^\dagger, L^\dagger}^{\pi^\dagger}(J^\dagger) = \pi \tau_{J^\dagger, \pi(L^\dagger)}(J^\dagger) = \pi(\pi(L^\dagger)) = L^\dagger \quad \text{and} \\ \pi^\dagger(\pi(J^\dagger)) &= \pi \alpha_{J^\dagger, L^\dagger}^{\pi^\dagger}(\pi(J^\dagger)) = \pi \tau_{J^\dagger, \pi(L^\dagger)} \tau_{L^\dagger, \pi(J^\dagger)} \pi(J^\dagger) = \pi(L^\dagger). \end{aligned}$$

Hence $\pi^\dagger$ is an involution on $\mathcal{I}$ and has cycles $(I^\dagger, K^\dagger), (J^\dagger, L^\dagger), (\pi(J^\dagger), \pi(L^\dagger))$.

In case ten, $|\mathcal{I}| = 8$, and $\pi^\dagger$ will be an involution on $\mathcal{I}$ with cycles $(I^\dagger, K^\dagger), (J^\dagger, L^\dagger), (\pi(I^\dagger), \pi(K^\dagger)), (\pi(J^\dagger), \pi(L^\dagger))$. As an illustration we note

$$\pi^\dagger(I^\dagger) = \pi \alpha_{I^\dagger, K^\dagger}^{\pi^\dagger} \alpha_{J^\dagger, L^\dagger}^{\pi^\dagger}(I^\dagger) = \pi \alpha_{I^\dagger, K^\dagger}^{\pi^\dagger}(I^\dagger) = \pi \tau_{I^\dagger, \pi(K^\dagger)} \tau_{K^\dagger, \pi(I^\dagger)}(I^\dagger) = \pi \tau_{I^\dagger, \pi(K^\dagger)}(I^\dagger) = \pi(\pi(K^\dagger)) = K^\dagger.$$

That $\pi^\dagger$ has the other cycles as claimed can be shown similarly. So in these two cases $\pi^\dagger$ is an involution restricted to $\mathcal{I}$ and has the cycles $(I^\dagger, K^\dagger), (J^\dagger, L^\dagger)$.

That $\pi^\dagger$ is an involution on $\mathcal{I}$ with cycles $(I^\dagger, K^\dagger), (J^\dagger, L^\dagger)$ can be similarly shown for the other cases, completing the proof. $\square$

Henceforth we will only write $\alpha_{i,j}$ for $\alpha_{i,j}^{\pi}$ unless otherwise mentioned. The utility of the construction (37) as a coupling is indicated by the following result.

Theorem 3.1. *Suppose π is chosen uniformly at random from Π_n and $(I^\dagger, J^\dagger, K^\dagger, L^\dagger)$ has joint distribution $p(\cdot)$ as in (36). If $\pi^\dagger$ is obtained from π according to (37) above, then π and $\pi^\dagger$ are constructed on a common space and $\pi^\dagger$ satisfies the conditions specified in Lemma 3.1.*

Proof. By hypothesis the indices $(I^\dagger, J^\dagger, K^\dagger, L^\dagger)$ have distribution in (36). From Lemma 3.2, we see that $\pi^\dagger$ is an involution and has cycles $(I^\dagger, K^\dagger), (J^\dagger, L^\dagger)$. It only remains to verify that the distribution of $\pi^\dagger$ is uniform over all involutions in Π_n having cycles $(I^\dagger, K^\dagger)$ and $(J^\dagger, L^\dagger)$. That is, recalling that $\mathbf{I}^\dagger = (I^\dagger, J^\dagger)$ and $\mathbf{K}^\dagger = (K^\dagger, L^\dagger)$, letting

$$\Pi_{n, \mathbf{I}^\dagger, \mathbf{K}^\dagger} = \{\pi \in \Pi_n : \pi(I^\dagger) = K^\dagger, \pi(J^\dagger) = L^\dagger\},$$

we need to verify that

$$P(\pi^\dagger = \phi | \mathbf{I}^\dagger, \mathbf{K}^\dagger) = \frac{1}{|\Pi_{n, \mathbf{I}^\dagger, \mathbf{K}^\dagger}|} = \frac{1}{|\Pi_{n-4}|} \quad \text{for all } \phi \in \Pi_{n, \mathbf{I}^\dagger, \mathbf{K}^\dagger}. \quad (38)$$

Since $I^\dagger, J^\dagger, K^\dagger, L^\dagger$ are distinct, with $\mathcal{I}$ as in Lemma 3.2, the size of $\mathcal{I}$ satisfies $4 \leq |\mathcal{I}| \leq 8$. In addition, since π is an involution we see that $|\{I^\dagger, J^\dagger, K^\dagger, L^\dagger\} \cap \{\pi(I^\dagger), \pi(J^\dagger), \pi(K^\dagger), \pi(L^\dagger)\}|$ is even. Hence so is $|\mathcal{I}|$, and we conclude that $\mathcal{I} \in \{4, 6, 8\}$.

For $\phi \in \Pi_{n, \mathbf{I}^\dagger, \mathbf{K}^\dagger}$, independence of $\mathcal{I}$ and $(\mathbf{I}^\dagger, \mathbf{K}^\dagger)$ yields

$$\begin{aligned} P(\pi^\dagger = \phi | \mathbf{I}^\dagger, \mathbf{K}^\dagger) &= \sum_{\iota \in \{4, 6, 8\}} P(\pi^\dagger = \phi | |\mathcal{I}| = \iota, \mathbf{I}^\dagger, \mathbf{K}^\dagger) P(|\mathcal{I}| = \iota | \mathbf{I}^\dagger, \mathbf{K}^\dagger) \\ &= \sum_{\iota \in \{4, 6, 8\}} P(\pi^\dagger = \phi | |\mathcal{I}| = \iota, \mathbf{I}^\dagger, \mathbf{K}^\dagger) P(|\mathcal{I}| = \iota). \end{aligned} \quad (39)$$

For $\pi \in \Pi_n$ let $\bar{\pi}$ denote the restriction of π to the complement of $I^\dagger, J^\dagger, K^\dagger, L^\dagger$. First consider the case $\iota = 4$ that is $\mathcal{I} = \{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}$. Since $\pi^\dagger \in \Pi_{n, \mathbf{I}^\dagger, \mathbf{K}^\dagger}$ the permutation $\pi^\dagger$ agrees with every $\phi \in \Pi_{n, \mathbf{I}^\dagger, \mathbf{K}^\dagger}$ on $I^\dagger, J^\dagger, K^\dagger, L^\dagger$, and, as $\pi^\dagger$ and π agree on $\mathcal{I}^c$,

$$\begin{aligned} P(\pi^\dagger = \phi | |\mathcal{I}| = 4, \mathbf{I}^\dagger, \mathbf{K}^\dagger) &= P(\bar{\pi}^\dagger = \bar{\phi} | |\mathcal{I}| = 4, \mathbf{I}^\dagger, \mathbf{K}^\dagger) \\ &= P(\bar{\pi} = \bar{\phi} | |\mathcal{I}| = 4, \mathbf{I}^\dagger, \mathbf{K}^\dagger) \\ &= \frac{1}{|\Pi_{n-4}|}. \end{aligned}$$

Now suppose $\iota = 6$. In this case, the set $\mathcal{J} = \mathcal{I} \setminus \{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}$ has size 2, say $\mathcal{J} = \{i_1, i_2\}$. We claim that $\pi^\dagger$ has the cycle (i_1, i_2) with $\{i_1, i_2\} = \{\pi(a), \pi(b)\}$ for some $a, b \in \{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}$, and that conditional on $|\mathcal{I}| = 6$ and $\{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}$ the values i_1, i_2 are uniform over all pairs of distinct values in $\{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}^c$. That $\pi^\dagger$ has the cycle (i_1, i_2) follows from Lemma 3.2. Suppose $(i_1, i_2) = (\pi(J^\dagger), \pi(L^\dagger))$ as in case 1 in (37), since $J^\dagger, L^\dagger$ do not form a cycle, and their images under π under case 1 is constrained exactly to lie outside $\{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}$, as π is uniform over Π_n , these images are uniform over $\{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}^c$. One can show that these properties hold similarly in the remaining cases. The remaining cycles of $\pi^\dagger$ are in $\mathcal{I}^c$ and thus are the same as those of $\pi|_{\mathcal{I}^c}$. Thus, the cycles of $\pi^\dagger$ are conditionally uniform, that is, for $\phi \in \Pi_{n, \mathbf{I}^\dagger, \mathbf{K}^\dagger}$

$$P(\pi^\dagger = \phi | |\mathcal{I}| = 6, \mathbf{I}^\dagger, \mathbf{K}^\dagger) = P(\bar{\pi}^\dagger = \bar{\phi} | |\mathcal{I}| = 6, \mathbf{I}^\dagger, \mathbf{K}^\dagger) = \frac{1}{|\Pi_{n-4}|}.$$

The case $\iota = 8$, that is, where $R_1 = R_2 = 0$, is handled similar to $\iota = 6$. Here $\pi^\dagger$ will have the cycles $(\pi(I^\dagger), \pi(K^\dagger)), (\pi(J^\dagger), \pi(L^\dagger))$. These two cycles are both of the form $(\pi(a), \pi(b))$ with $a, b \in \{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}$ and hence, as in the case $\iota = 6$, uniform random transpositions. Since, π and $\pi^\dagger$ agree on $\mathcal{I}^c$, we can see that $\pi^\dagger$ has uniform random transpositions on $\{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}^c = \mathcal{I}^c \cup \{\pi(I^\dagger), \pi(J^\dagger), \pi(K^\dagger), \pi(L^\dagger)\}$ yielding

$$P(\pi^\dagger = \phi | |\mathcal{I}| = 8, \mathbf{I}^\dagger, \mathbf{K}^\dagger) = P(\bar{\pi}^\dagger = \bar{\phi} | |\mathcal{I}| = 8, \mathbf{I}^\dagger, \mathbf{K}^\dagger) = \frac{1}{|\Pi_{n-4}|}.$$

Thus (39) now yields

$$P(\pi^\dagger = \phi) = \frac{1}{|\Pi_{n-4}|},$$

verifying (38) and proving the theorem. $\square$

So the tuple $(\pi^\dagger, \pi^\ddagger)$ obtained in Theorem 3.1 satisfy the conditions in Lemma 3.1. Hence $(W^\dagger, W^\ddagger)$ constructed from $(\pi^\dagger, \pi^\ddagger)$ as in Lemma 3.1 has the required square bias distribution.

We conclude this section with the calculation of the normalization constant for the distribution $p(\cdot)$ in (36).

Lemma 3.3. *For $n \geq 4$ and $D = ((d_{ij}))_{n \times n}$ satisfying (15), we have*

$$c_n \sum_{|\{i, j, k, l\}|=4} [d_{ik} + d_{jl} - (d_{ij} + d_{kl})]^2 = 1 \quad \text{where} \quad c_n = \frac{1}{2(n-1)^2(n-3)} = O\left(\frac{1}{n^3}\right), \quad (40)$$

and in particular

$$c_n \leq \frac{1}{n^3} \quad \text{when } n \geq 9. \quad (41)$$

Proof. From (28), we have

$$W - W' = 2(d_{I\pi(I)} + d_{J\pi(J)} - (d_{IJ} + d_{\pi(I)\pi(J)})),$$

where (I, J) are two distinct indices selected uniformly from $\{1, 2, \dots, n\}$. Since π is an involution chosen uniformly, we have

$$\begin{aligned} E(W - W')^2 &= \frac{1}{n(n-1)} \sum_{i \neq j} 4E(d_{i\pi(i)} + d_{j\pi(j)} - (d_{ij} + d_{\pi(i)\pi(j)}))^2 \\ &= \frac{4}{n(n-1)^2(n-3)} \sum_{|\{i,j,k,l\}|=4} (d_{ik} + d_{jl} - (d_{ij} + d_{kl}))^2. \end{aligned} \quad (42)$$

Using (42), (17) and (29) and $\sigma_D^2 = 1$, we obtain

$$\frac{4}{n(n-1)^2(n-3)} \sum_{|\{i,j,k,l\}|=4} (d_{ik} + d_{jl} - (d_{ij} + d_{kl}))^2 = 2\lambda\sigma_D^2 = \frac{8}{n}.$$

On simplification, we obtain

$$\frac{1}{2(n-1)^2(n-3)} \sum_{|\{i,j,k,l\}|=4} (d_{ik} + d_{jl} - (d_{ij} + d_{kl}))^2 = 1,$$

proving (40). The verification of (41) is direct. $\square$

4 L^1 bounds

In this section we derive the L^1 bounds for the normal approximation of $W = \sum_{i=1}^n d_{i\pi(i)}$, where π is chosen uniformly at random from Π_n . The main theorem in this section is the following.

Theorem 4.1. *Let π be an involution chosen uniformly at random from Π_n and $D = ((d_{ij}))$ be an array satisfying (15). Then with β_D as in (11), $W = \sum_{i=1}^n d_{i\pi(i)}$ satisfies*

$$\|F_W - \Phi\|_1 \leq \frac{\beta_D}{n} \left(224 + 1344\frac{1}{n} + 384\frac{1}{n^2} \right) \quad \text{when } n \geq 9.$$

In particular,

$$\|F_W - \Phi\|_1 \leq 379\frac{\beta_D}{n} \quad \text{when } n \geq 9.$$

We will need the following inequalities in order to prove Theorem 4.1. To avoid writing down the indices over which we are summing, unless otherwise specified the summation will be taken over the same index set as the one immediately preceding it.

With $p(\cdot)$ as in (36), in what follows, we will apply bounds such as

$$\begin{aligned} \sum_{|\{i,j,k,l\}|=4} |d_{ik}|p(i,j,k,l) &= c_n \sum |d_{ik}|[d_{ik} + d_{jl} - (d_{ij} + d_{kl})]^2 \\ &\leq \sum_{i,j,k,l} |d_{ik}|[d_{ik} + d_{jl} - (d_{ij} + d_{kl})]^2 \\ &= c_n \sum |d_{ik}|(d_{ik}^2 + d_{jl}^2 + d_{ij}^2 + d_{kl}^2) \\ &\leq 4c_n n^2 \beta_D \leq 4\frac{\beta_D}{n} \quad \text{when } n \geq 9, \text{ from (41)}. \end{aligned} \quad (43)$$

The first nontrivial equality above uses the special form of the term inside squares. Whenever we encounter a cross term we always get a free index to sum over which gives us zero since $d_{i+} = 0 \forall i$. The second inequality uses the fact that for any choices $\iota_1, \iota_2, \kappa_1, \kappa_2 \in \{i, j, k, l\}$ with $\iota_1 \neq \kappa_1$ and $\iota_2 \neq \kappa_2$, perhaps by relabelling the indices after the inequality,

$$\sum_{i,j,k,l} |d_{\iota_1 \kappa_1}| d_{\iota_2 \kappa_2}^2 \leq (\sum |d_{ij}|^3)^{\frac{1}{3}} (\sum |d_{kl}|^3)^{\frac{2}{3}} \leq n^2 \beta_D.$$

Generally the exponent of n in such an inequality will be 2 less the number of indices over which we are summing up. For instance, if we are summing up over 5 indices the exponent of n will be 3 and so on. In particular,

$$\begin{aligned} \sum_{|\{i,j,k,l,s\}|=5} |d_{\iota s}| p(i, j, k, l) &\leq 4n^3 c_n \beta_D \quad \text{where } \iota \in \{i, j, k, l\} \\ &\leq 4\beta_D \quad \text{when } n \geq 9, \text{ using (41) and} \end{aligned} \quad (44)$$

$$\sum_{|\{i,j,k,l,s,t\}|=6} |d_{st}| p(i, j, k, l) \leq 4n^4 c_n \beta_D \leq 4n\beta_D \quad \text{when } n \geq 9. \quad (45)$$

Theorem 4.2. Suppose $D = ((d_{ij}))$ is an array satisfying (15) and π and $\pi^\dagger$ are as in Theorem 3.1, and $\pi^\ddagger, W^\dagger, W^\ddagger$ and W^* are as in Lemma 3.1. Then $W, W^\dagger, W^\ddagger$ can be decomposed as

$$W = S + T \quad W^\dagger = S + T^\dagger \quad W^\ddagger = S + T^\ddagger, \quad (46)$$

where

$$S = \sum_{i \notin \mathcal{I}} d_{i\pi(i)}, \quad T = \sum_{i \in \mathcal{I}} d_{i\pi(i)}, \quad T^\dagger = \sum_{i \in \mathcal{I}} d_{i\pi^\dagger(i)} \quad \text{and} \quad T^\ddagger = \sum_{i \in \mathcal{I}} d_{i\pi^\ddagger(i)}, \quad (47)$$

where $\mathcal{I}$ is as in Lemma 3.2. Also, W^* has the W zero bias distribution and satisfies

$$E|W - W^*| \leq 112 \frac{\beta_D}{n} + 672 \frac{\beta_D}{n^2} + 192 \frac{\beta_D}{n^3} \quad \text{when } n \geq 9. \quad (48)$$

In view of (13) Theorem 4.2 implies Theorem 4.1.

Proof. Lemma 3.1 guarantees that W^* has the W -zero biased distribution. Recalling $\mathcal{I} = \{I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger), \pi(K^\dagger), \pi(L^\dagger)\}$ and that $\pi, \pi^\dagger$ and $\pi^\ddagger$ agree on $\mathcal{I}^c$ by Lemma 3.2, we obtain decomposition (46).

From $W^* = UW^\dagger + (1 - U)W^\ddagger$ and (46), we obtain

$$E|W - W^*| = E|UT^\dagger + (1 - U)T^\ddagger - T|. \quad (49)$$

Using the fact that $E(U) = 1/2$, and that U is independent of $T^\dagger$ and $T^\ddagger$, we obtain

$$E|W - W^*| \leq \frac{1}{2}(E|T^\dagger| + E|T^\ddagger|) + E|T| = EV \quad \text{where } V = |T^\dagger| + |T|, \quad (50)$$

where the equality follows from the fact that $\pi^\dagger, \pi^\ddagger$, and therefore $T^\dagger, T^\ddagger$, are exchangeable.

Thus our goal is to bound the L^1 norms of T and $T^\dagger$ and we proceed in a case by case basis, much along the lines of Section 6 in [3]. In summary, we group the ten cases in (37) into the following five cases: $R_1 = 1$; $R_1 = 0, R_2 = 1$; $R_1 = 2$; $R_1 = 0, R_2 = 2$; $R_1 = 0, R_2 = 0$.

Computation on $R_1 = 1$: The event $R_1 = 1$, which we indicate by $\mathbf{1}_1$, can occur in four different ways, corresponding to the first four cases in the definition of $\pi^\ddagger$ in (37). With V as in (50), we can decompose $\mathbf{1}_1$ to yield

$$V\mathbf{1}_1 = V\mathbf{1}_{1,1} + V\mathbf{1}_{1,2} + V\mathbf{1}_{1,3} + V\mathbf{1}_{1,4}, \quad (51)$$

where $\mathbf{1}_{1,1} = \mathbf{1}(\pi(I^\dagger) = K^\dagger, \pi(J^\dagger) \neq L^\dagger)$ and $\mathbf{1}_{1,m}$ for $m = 2, 3, 4$ similarly corresponding to the other three cases in (37), in their respective order. On $\mathbf{1}_{1,1}$, we have $\mathcal{I} = \{I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(J^\dagger), \pi(L^\dagger)\}$ and $\pi(I^\dagger) = K^\dagger$, yielding

$$T\mathbf{1}_{1,1} = \sum_{i \in \mathcal{I}} d_{i\pi(i)} \mathbf{1}_{1,1} = 2(d_{I^\dagger K^\dagger} + d_{J^\dagger \pi(J^\dagger)} + d_{L^\dagger \pi(L^\dagger)}) \mathbf{1}_{1,1}. \quad (52)$$

By Lemma 3.2 $\pi^\dagger$ has cycles $(I^\dagger, K^\dagger), (J^\dagger, L^\dagger)$ and is an involution restricted to $\mathcal{I}$, hence

$$T^\dagger \mathbf{1}_{1,1} = \sum_{i \in \mathcal{I}} d_{i\pi^\dagger(i)} \mathbf{1}_{1,1} = 2(d_{I^\dagger K^\dagger} + d_{J^\dagger L^\dagger} + d_{\pi(J^\dagger)\pi(L^\dagger)}) \mathbf{1}_{1,1}. \quad (53)$$

So, we obtain

$$E|T|\mathbf{1}_{1,1} \leq 2(E|d_{I^\dagger K^\dagger}|\mathbf{1}_{1,1} + E|d_{J^\dagger \pi(J^\dagger)}|\mathbf{1}_{1,1} + E|d_{L^\dagger \pi(L^\dagger)}|\mathbf{1}_{1,1}) \quad (54)$$

$$E|T^\dagger|\mathbf{1}_{1,1} \leq 2(E|d_{I^\dagger, K^\dagger}|\mathbf{1}_{1,1} + E|d_{J^\dagger, L^\dagger}|\mathbf{1}_{1,1} + E|d_{\pi(J^\dagger), \pi(L^\dagger)}|\mathbf{1}_{1,1})). \quad (55)$$

Because of the indicator in (54) and (55), we need to consider the joint distribution

$$p_2(i, j, k, l, s, t) = P((I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger)) = (i, j, k, l, s, t)),$$

which includes the images of $I^\dagger, J^\dagger$ under π , say s and t respectively. With c_n as in Lemma 3.3, we claim $p_2(\cdot)$ is given by

$$p_2(i, j, k, l, s, t) = \begin{cases} \frac{1}{n-1}p(i, j, k, l) & \text{when } s = j, t = i \\ 0 & \text{when } s = j, t \neq i \text{ or } s \neq j, t = i \\ 0 & \text{when } s = i \text{ or } t = j \\ 0 & \text{when } s = t \\ \frac{1}{(n-1)(n-3)}p(i, j, k, l) & \text{when } t \notin \{j, s, i\} \Leftrightarrow s \notin \{i, t, j\}. \end{cases} \quad (56)$$

To justify (56), note first that $s = j$ if and only if $t = i$, for example, $s = j$ implies $t = \pi(j) = \pi(s) = \pi(\pi(i)) = i$, and therefore

$$\{s = j\} = \{s = j, t = i\} = \{t = i\}. \quad (57)$$

Thus the second case of (56) has zero probability. The remaining trivial cases can be discarded using the fact that $\pi \in \Pi_n$. Leaving these out, the first probability is derived using (57), since the image of $I^\dagger$ under π is uniform over $\{I^\dagger\}^c$ and independent of $(I^\dagger, J^\dagger, K^\dagger, L^\dagger)$, so in particular takes the value $s = j$ with probability $1/(n-1)$. In the last case it is easy to see that $t \notin \{j, s, i\}$ and $s \notin \{i, t, j\}$ are equivalent. The image of $I^\dagger$ is uniform over all available $n-1$ choices, and, conditional on $\pi(I^\dagger) \neq J^\dagger$, the $n-3$ remaining choices for the image of $J^\dagger$ fall in $\{I^\dagger, \pi(I^\dagger), J^\dagger\}^c$ uniformly.

Next we bound each of the summands in (54) and (55) separately. First note that under $\mathbf{1}_{1,1}$ only the last form of $p_2(i, j, k, l, s, t)$ in (56) is relevant. In particular, $s \neq t$ always, and since i, j, k, l are distinct, $s = k$ implies $s \notin \{i, j\}$. Hence, for the first summand in (54), we obtain

$$\begin{aligned} E|d_{I^\dagger K^\dagger}|\mathbf{1}_{1,1} &= \sum_{i, j, k, l, s, t} |d_{ik}| p_2(i, j, k, l, s, t) \mathbf{1}(s = k, t \neq l) \\ &= \sum_{|\{i, j, k, l, t\}|=5} |d_{ik}| p_2(i, j, k, l, k, t) \\ &= \frac{n-4}{(n-1)(n-3)} \sum_{|\{i, j, k, l\}|=4} |d_{ik}| p(i, j, k, l) \\ &\leq \frac{4}{n(n-1)} \beta_D \quad \text{using (43)} \\ &\leq 8 \frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \end{aligned} \quad (58)$$

Similarly, we can estimate the second summand in (54) as

$$\begin{aligned}
E|d_{J^\dagger \pi(J^\dagger)}| \mathbf{1}_{1,1} &= \sum_{i,j,k,l,s,t} |d_{jt}| p_2(i,j,k,l,s,t) \mathbf{1}(s=k, t \neq l) \\
&= \sum_{|\{i,j,k,l,t\}|=5} |d_{jt}| p_2(i,j,k,l,k,t) \\
&= \frac{1}{(n-1)(n-3)} \sum |d_{jt}| p(i,j,k,l) \\
&\leq \frac{4}{(n-1)(n-3)} \beta_D \quad \text{using (44), when } n \geq 9 \\
&\leq 8 \frac{\beta_D}{n^2} \quad \text{when } n \geq 9.
\end{aligned} \tag{59}$$

The last summand in (54) is $E|d_{L^\dagger \pi(L^\dagger)}| \mathbf{1}_{1,1}$. Using the fact that

$$(I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger)) \stackrel{\mathcal{L}}{=} (K^\dagger, L^\dagger, I^\dagger, J^\dagger, \pi(K^\dagger), \pi(L^\dagger)),$$

where $\stackrel{\mathcal{L}}{=}$ denotes equality in distribution, (59) yields,

$$E|d_{L^\dagger \pi(L^\dagger)}| \mathbf{1}_{1,1} = E|d_{J^\dagger \pi(J^\dagger)}| \mathbf{1}_{1,1} \leq 8 \frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \tag{60}$$

Thus combining the bounds in (58),(59),(60) we obtain the following bound on the term (54),

$$E|T| \mathbf{1}_{1,1} \leq 48 \frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \tag{61}$$

Now moving to (55), we note the first summand in (55) is the same as the first summand of (54), and so can be bounded by (58). We can bound $E|d_{J^\dagger L^\dagger}| \mathbf{1}_{1,1}$, which is the second summand in (55) in a similar fashion as in (58) through the following calculation

$$\begin{aligned}
E|d_{J^\dagger L^\dagger}| \mathbf{1}_{1,1} &= \sum_{|\{i,j,k,l,t\}|=5} |d_{jl}| p_2(i,j,k,l,k,t) \\
&= \frac{n-4}{(n-1)(n-3)} \sum_{|\{i,j,k,l\}|=4} |d_{jl}| p(i,j,k,l) \\
&\leq 8 \frac{\beta_D}{n^2} \quad \text{when } n \geq 9, \text{ using (43)}.
\end{aligned} \tag{62}$$

So we are only left with the last summand of (55) which is $E|d_{\pi(J^\dagger) \pi(L^\dagger)}| \mathbf{1}_{1,1}$. For this we will need to introduce the joint distribution

$$p_3(i,j,k,l,s,t,r) = P((I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger), \pi(L^\dagger)) = (i,j,k,l,s,t,r)). \tag{63}$$

The case $\mathbf{1}_{1,1}$ is equivalent to $s=k, t \neq l, r \neq j$, and, since $\pi \in \Pi_n$ it is equivalent to $s=k$ and $|\{i,j,k,l,r,t\}|=6$. We claim that

$$p_3(i,j,k,l,s,t,r) = \frac{1}{(n-1)(n-3)(n-5)} p(i,j,k,l) \quad \text{when } s=k \text{ and } |\{i,j,k,l,r,t\}|=6. \tag{64}$$

To justify (64), we note that since π is independent of $\{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}$, the image of $I^\dagger$ is uniform over $n-1$ choices from $\{I^\dagger\}^c$ and conditional on $\pi(I^\dagger) = K^\dagger$, the $n-3$ choices for $\pi(J^\dagger)$ fall in $\{I^\dagger, J^\dagger, K^\dagger\}^c$ uniformly. Conditioned on these two images, when $\pi(J^\dagger) \neq L^\dagger$, $\pi(L^\dagger)$ is distributed uniformly over the $n-5$ choices of $\{I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(J^\dagger)\}^c$.

Now we can bound $E|d_{\pi(J^\dagger)\pi(L^\dagger)}|\mathbf{1}_{1,1}$ in the following way,

$$\begin{aligned}
E|d_{\pi(J^\dagger)\pi(L^\dagger)}|\mathbf{1}_{1,1} &= \sum_{i,j,k,l,s,t,r} |d_{tr}|p_3(i,j,k,l,s,t,r)\mathbf{1}(s=k, t \neq l, r \neq j) \\
&= \sum_{|\{i,j,k,l,t,r\}|=6} |d_{tr}|p_3(i,j,k,l,k,t,r) \\
&= \frac{1}{(n-1)(n-3)(n-5)} \sum |d_{tr}|p(i,j,k,l) \\
&\leq \frac{4n}{(n-1)(n-3)(n-5)}\beta_D \quad \text{using (45)} \\
&\leq 16\frac{\beta_D}{n^2} \quad \text{when } n \geq 9.
\end{aligned} \tag{65}$$

So, adding the bounds in (58),(62) and (65), and using (55) we obtain,

$$E|T^\dagger|\mathbf{1}_{1,1} \leq 64\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \tag{66}$$

From (61) and (66), using the definition of V in (50), we obtain the following bound on the first term of (51),

$$EV\mathbf{1}_{1,1} \leq 112\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \tag{67}$$

Next, on $\mathbf{1}_{1,2}$, indicating the event $\pi(I^\dagger) \neq K^\dagger, \pi(J^\dagger) = L^\dagger$, we have $\mathcal{I} = \{I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(K^\dagger)\}$ and hence, by definition (47),

$$T\mathbf{1}_{1,2} = 2(d_{I^\dagger\pi(I^\dagger)} + d_{K^\dagger\pi(K^\dagger)} + d_{J^\dagger L^\dagger})\mathbf{1}_{1,2} \tag{68}$$

We further observe,

$$(I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger), \pi(K^\dagger), \pi(L^\dagger)) \stackrel{\mathcal{L}}{=} (J^\dagger, I^\dagger, L^\dagger, K^\dagger, \pi(J^\dagger), \pi(I^\dagger), \pi(L^\dagger), \pi(K^\dagger)), \tag{69}$$

which because of (61) yields

$$T\mathbf{1}_{1,2} \stackrel{\mathcal{L}}{=} T\mathbf{1}_{1,1} \Rightarrow E|T|\mathbf{1}_{1,2} = E|T|\mathbf{1}_{1,1} \leq 48\frac{\beta_D}{n^2} \quad \text{for } n \geq 9 \tag{70}$$

Furthermore, the distributional equality in (69) implies

$$T^\dagger\mathbf{1}_{1,2} = 2(d_{I^\dagger K^\dagger} + d_{J^\dagger L^\dagger} + d_{\pi(I^\dagger)\pi(K^\dagger)})\mathbf{1}_{1,2} \stackrel{\mathcal{L}}{=} T^\dagger\mathbf{1}_{1,1}, \tag{71}$$

yielding

$$E|T^\dagger|\mathbf{1}_{1,2} = E|T^\dagger|\mathbf{1}_{1,1} \leq 64\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \tag{72}$$

Thus combining (70),(72) we obtain

$$EV\mathbf{1}_{1,2} \leq 112\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \tag{73}$$

Next on $\mathbf{1}_{1,3}$, indicating $\pi(I^\dagger) = L^\dagger, \pi(J^\dagger) \neq K^\dagger$, we have $\mathcal{I} = \{I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(J^\dagger), \pi(K^\dagger)\}$ and

$$T\mathbf{1}_{1,3} = 2(d_{I^\dagger L^\dagger} + d_{J^\dagger\pi(J^\dagger)} + d_{K^\dagger\pi(K^\dagger)})\mathbf{1}_{1,3}, \tag{74}$$

$$T^\dagger\mathbf{1}_{1,3} = 2(d_{I^\dagger K^\dagger} + d_{J^\dagger L^\dagger} + d_{\pi(J^\dagger)\pi(K^\dagger)})\mathbf{1}_{1,3}. \tag{75}$$

On $\mathbf{1}_{1,3}$, we have $s = l, t \neq k$ which is equivalent to $s = l$ and $|\{i, j, k, l, t\}| = 5$. Hence we may bound the first summand in (74) as follows

$$\begin{aligned}
E|d_{I^\dagger L^\dagger}|\mathbf{1}_{1,3} &= \sum_{i,j,k,l,s,t} |d_{il}|p_2(i, j, k, l, s, t)\mathbf{1}(s = l, t \neq k) \\
&= \sum_{|\{i,j,k,l,t\}|=5} |d_{il}|p_2(i, j, k, l, l, t) \\
&= \frac{n-4}{(n-1)(n-3)} \sum_{|\{i,j,k,l\}|=4} |d_{il}|p(i, j, k, l) \\
&\leq \frac{4}{n(n-1)}\beta_D \quad \text{using (43)} \\
&\leq 8\frac{\beta_D}{n^2} \quad \text{when } n \geq 9.
\end{aligned}$$

Continuing in this manner we arrive, as in (73), at

$$EV\mathbf{1}_{1,3} \leq 112\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \quad (76)$$

Symmetries between $\mathbf{1}_{1,1}$ and $\mathbf{1}_{1,2}$ such as (69) hold as well between $\mathbf{1}_{1,3}$ and $\mathbf{1}_{1,4}$, yielding

$$EV\mathbf{1}_{1,4} \leq 112\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \quad (77)$$

Combining the bounds from (67), (73), (76) and (77), we obtain

$$EV\mathbf{1}_1 \leq 448\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \quad (78)$$

Computation on $R_1 = 0, R_2 = 1$: Now we need to make the following decomposition

$$V\mathbf{1}_2 = V\mathbf{1}_{2,1} + V\mathbf{1}_{2,2},$$

where $\mathbf{1}_2 = \mathbf{1}(R_1 = 0, R_2 = 1)$, $\mathbf{1}_{2,1} = \mathbf{1}(\pi(I^\dagger) = J^\dagger, \pi(K^\dagger) \neq L^\dagger)$, $\mathbf{1}_{2,2} = \mathbf{1}(\pi(I^\dagger) \neq J^\dagger, \pi(K^\dagger) = L^\dagger)$. On $\mathbf{1}_{2,1}$ we have $\mathcal{I} = \{I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(K^\dagger), \pi(L^\dagger)\}$ which gives

$$\begin{aligned}
T\mathbf{1}_{2,1} &= 2(d_{I^\dagger J^\dagger} + d_{K^\dagger \pi(K^\dagger)} + d_{L^\dagger \pi(L^\dagger)})\mathbf{1}_{2,1} \quad \text{and} \\
T^\dagger\mathbf{1}_{2,1} &= 2(d_{I^\dagger K^\dagger} + d_{J^\dagger L^\dagger} + d_{\pi(K^\dagger)\pi(L^\dagger)})\mathbf{1}_{2,1}.
\end{aligned}$$

So, we obtain,

$$E|T|\mathbf{1}_{2,1} \leq 2(E|d_{I^\dagger J^\dagger}|\mathbf{1}_{2,1} + E|d_{K^\dagger \pi(K^\dagger)}|\mathbf{1}_{2,1} + E|d_{L^\dagger \pi(L^\dagger)}|\mathbf{1}_{2,1}), \quad (79)$$

$$E|T^\dagger|\mathbf{1}_{2,1} \leq 2(E|d_{I^\dagger K^\dagger}|\mathbf{1}_{2,1} + E|d_{J^\dagger L^\dagger}|\mathbf{1}_{2,1} + E|d_{\pi(K^\dagger)\pi(L^\dagger)}|\mathbf{1}_{2,1}). \quad (80)$$

Since $p(i, j, k, l) = p(i, k, j, l)$ and π is chosen independently of $\{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}$, we have

$$(I^\dagger, K^\dagger, J^\dagger, L^\dagger, \pi(I^\dagger), \pi(K^\dagger), \pi(J^\dagger), \pi(L^\dagger)) \stackrel{\mathcal{L}}{=} (I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger), \pi(K^\dagger), \pi(L^\dagger)). \quad (81)$$

Hence we obtain

$$T\mathbf{1}_{2,1} \stackrel{\mathcal{L}}{=} T\mathbf{1}_{1,1},$$

which, by (61) gives

$$E|T|\mathbf{1}_{2,1} = E|T|\mathbf{1}_{1,1} \leq 48\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \quad (82)$$

To begin bounding $E|T^\dagger|\mathbf{1}_{2,1}$, we bound the first summand in (80), $E|d_{I^\dagger K^\dagger}|\mathbf{1}_{2,1}$, as follows

$$\begin{aligned}
E|d_{I^\dagger K^\dagger}|\mathbf{1}_{2,1} &= \sum_{|\{i,j,k,l,r\}|=5} |d_{ik}|P((I^\dagger, K^\dagger, J^\dagger, L^\dagger, \pi(I^\dagger), \pi(K^\dagger)) = (i, k, j, l, j, r)) \\
&= \sum |d_{ik}|P((I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger)) = (i, k, j, l, j, r)) \quad \text{using (81),} \\
&= \sum |d_{ik}|p_2(i, k, j, l, j, r) \\
&= \frac{n-4}{(n-1)(n-3)} \sum_{|\{i,j,k,l\}|=4} |d_{ik}|p(i, j, k, l) \\
&\leq 8\frac{\beta_D}{n^2} \quad \text{when } n \geq 9, \text{ using (43).}
\end{aligned} \tag{83}$$

Also using the distributional equality $(I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(K^\dagger)) \stackrel{\mathcal{L}}{=} (J^\dagger, I^\dagger, L^\dagger, K^\dagger, \pi(J^\dagger), \pi(L^\dagger))$ and the bound in (83) above, we obtain

$$E|d_{J^\dagger L^\dagger}|\mathbf{1}_{2,1} = E|d_{I^\dagger K^\dagger}|\mathbf{1}_{2,1} \leq 8\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \tag{84}$$

Using the distributional equality in (81) and the bound in (65)

$$E|d_{\pi(K^\dagger)\pi(L^\dagger)}|\mathbf{1}_{2,1} = E|d_{\pi(J^\dagger)\pi(L^\dagger)}|\mathbf{1}_{1,1} \leq 16\frac{\beta_D}{n^2}. \tag{85}$$

Combining (84),(85) and using (80) we obtain

$$E|T^\dagger|\mathbf{1}_{2,1} \leq 64\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \tag{86}$$

Adding the two bounds in (82) and (86) we obtain

$$EV\mathbf{1}_{2,1} \leq 112\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \tag{87}$$

Next, on $\mathbf{1}_{2,2}$, where $\pi(I^\dagger) \neq J^\dagger, \pi(K^\dagger) = L^\dagger$, we have $\mathcal{I} = \{I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger)\}$ and hence

$$T\mathbf{1}_{2,2} = 2(d_{I^\dagger \pi(I^\dagger)} + d_{J^\dagger \pi(J^\dagger)} + d_{K^\dagger L^\dagger})\mathbf{1}_{2,2} \tag{88}$$

$$T^\dagger\mathbf{1}_{2,2} = 2(d_{I^\dagger K^\dagger} + d_{J^\dagger L^\dagger} + d_{\pi(I^\dagger)\pi(J^\dagger)})\mathbf{1}_{2,2}. \tag{89}$$

Noting the distributional equality

$$(I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger), \pi(K^\dagger), \pi(L^\dagger)) \stackrel{\mathcal{L}}{=} (K^\dagger, L^\dagger, I^\dagger, J^\dagger, \pi(K^\dagger), \pi(L^\dagger), \pi(I^\dagger), \pi(J^\dagger))$$

we obtain

$$T\mathbf{1}_{2,1} \stackrel{\mathcal{L}}{=} T\mathbf{1}_{2,2} \quad \text{and} \quad T^\dagger\mathbf{1}_{2,1} \stackrel{\mathcal{L}}{=} T^\dagger\mathbf{1}_{2,2},$$

which yields

$$E|T|\mathbf{1}_{2,2} = E|T|\mathbf{1}_{2,1} \leq 48\frac{\beta_D}{n^2} \quad \text{and} \quad E|T^\dagger|\mathbf{1}_{2,2} = E|T^\dagger|\mathbf{1}_{2,1} \leq 64\frac{\beta_D}{n^2}.$$

Hence we have

$$EV\mathbf{1}_{2,2} \leq 112\frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \tag{90}$$

Combining (87) with (90) we obtain

$$EV\mathbf{1}_2 \leq 224 \frac{\beta_D}{n^2} \quad \text{when } n \geq 9. \quad (91)$$

Computation on $R_1 = 2$: Here we need the decomposition

$$V\mathbf{1}_3 = V\mathbf{1}_{3,1} + V\mathbf{1}_{3,2},$$

where $\mathbf{1}_3 = \mathbf{1}(R_1 = 2)$, $\mathbf{1}_{3,1} = \mathbf{1}(\pi(I^\dagger) = K^\dagger, \pi(J^\dagger) = L^\dagger)$ and $\mathbf{1}_{3,2} = \mathbf{1}(\pi(I^\dagger) = L^\dagger, \pi(J^\dagger) = K^\dagger)$. Note that $\mathbf{1}_{3,1}$ and $\mathbf{1}_{3,2}$ correspond to the cases in (37) where $(R_1, R_2) = (2, 0)$ and $(R_1, R_2) = (2, 2)$, respectively. On both $\mathbf{1}_{3,1}$ and $\mathbf{1}_{3,2}$, we have $\mathcal{I} = \{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}$ and

$$T^\dagger \mathbf{1}_{3,1} = T\mathbf{1}_{3,1} = 2(d_{I^\dagger K^\dagger} + d_{J^\dagger L^\dagger})\mathbf{1}_{3,1} \quad \text{since } \pi\mathbf{1}_{3,1} = \pi^\dagger \mathbf{1}_{3,1}. \quad (92)$$

From $(I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger)) \stackrel{\mathcal{L}}{=} (J^\dagger, I^\dagger, L^\dagger, K^\dagger, \pi(J^\dagger), \pi(I^\dagger))$, it is clear that $E|d_{I^\dagger K^\dagger}|\mathbf{1}_{3,1} = E|d_{J^\dagger L^\dagger}|\mathbf{1}_{3,1}$ and hence it is enough to bound any one of the two summands in (92). We bound $E|d_{I^\dagger K^\dagger}|\mathbf{1}_{3,1}$ using (43) as follows

$$\begin{aligned} E|d_{I^\dagger K^\dagger}|\mathbf{1}_{3,1} &= \sum_{i,j,k,l,s,t} |d_{ik}|p_2(i,j,k,l,s,t)\mathbf{1}(s=k, t=l) \\ &= \sum_{|\{i,j,k,l\}|=4} |d_{ik}|p_2(i,j,k,l,k,l) \\ &= \frac{1}{(n-1)(n-3)} \sum |d_{ik}|p(i,j,k,l) \\ &\leq 8 \frac{\beta_D}{n^3} \quad \text{when } n \geq 9. \end{aligned} \quad (93)$$

Thus, using (92) and (93), we obtain

$$EV\mathbf{1}_{3,1} \leq 64 \frac{\beta_D}{n^3} \quad \text{when } n \geq 9. \quad (94)$$

On $\mathbf{1}_{3,2}$, we have

$$T\mathbf{1}_{3,2} = 2(d_{I^\dagger L^\dagger} + d_{J^\dagger K^\dagger}) \quad \text{and} \quad T^\dagger \mathbf{1}_{3,2} = 2(d_{I^\dagger K^\dagger} + d_{J^\dagger L^\dagger}). \quad (95)$$

To obtain bounds for $E(V\mathbf{1}_{3,2})$, we bound the first summand in (95) as follows,

$$\begin{aligned} E|d_{I^\dagger L^\dagger}|\mathbf{1}_{3,2} &= \sum_{|\{i,j,k,l\}|=4} |d_{il}|p_2(i,j,k,l,l,k) \\ &= \frac{1}{(n-1)(n-3)} \sum |d_{il}|p(i,j,k,l) \leq 8 \frac{\beta_D}{n^3} \quad \text{when } n \geq 9, \text{ using (43)}. \end{aligned} \quad (96)$$

Similarly its easy to obtain bounds on the other summands also and conclude as in (94),

$$EV\mathbf{1}_{3,2} \leq 64 \frac{\beta_D}{n^3} \quad \text{for } n \geq 9. \quad (97)$$

Combining (94) and (97), we obtain

$$EV\mathbf{1}_3 \leq 128 \frac{\beta_D}{n^3} \quad \text{when } n \geq 9. \quad (98)$$

Computation for $R_1 = 0, R_2 = 2$: This event is indicated by $\mathbf{1}_4 = \mathbf{1}(\pi(I^\dagger) = J^\dagger, \pi(K^\dagger) = L^\dagger)$. On $\mathbf{1}_4$, we have

$$T\mathbf{1}_4 = 2(d_{I^\dagger J^\dagger} + d_{K^\dagger L^\dagger})\mathbf{1}_4, \quad (99)$$

$$T^\dagger \mathbf{1}_4 = 2(d_{I^\dagger K^\dagger} + d_{J^\dagger L^\dagger})\mathbf{1}_4. \quad (100)$$

For the first term in (99),

$$\begin{aligned}
E|d_{I^\dagger J^\dagger}|\mathbf{1}_4 &= \sum_{|\{i,j,k,l\}|=4} |d_{ij}| P((I^\dagger, K^\dagger, J^\dagger, L^\dagger, \pi(I^\dagger), \pi(K^\dagger)) = (i, k, j, l, j, l)) \\
&= \sum |d_{ij}| p_2(i, k, j, l, j, l) \quad \text{using (81)} \\
&\leq \frac{1}{(n-1)(n-3)} \sum |d_{ij}| p(i, j, k, l) \\
&\leq 8 \frac{\beta_D}{n^3} \quad \text{when } n \geq 9.
\end{aligned} \tag{101}$$

For the other summand in (99) and (100), we can follow similar calculations as in (101) above and obtain the same bounds. Thus we will finally obtain

$$E|T\mathbf{1}_4|, E|T^\dagger\mathbf{1}_4| \leq 32 \frac{\beta_D}{n^3} \quad \text{when } n \geq 9. \tag{102}$$

So we have

$$EV\mathbf{1}_4 \leq 64 \frac{\beta_D}{n^3} \quad \text{when } n \geq 9. \tag{103}$$

Computation on $R_1 = R_2 = 0$: Now we bound the L^1 contribution from the last case in (37) denoted by $\mathbf{1}_5 = \mathbf{1}(R_1 = R_2 = 0)$. Here we have $\mathcal{I} = \{I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger), \pi(K^\dagger), \pi(L^\dagger)\}$ and

$$T\mathbf{1}_5 = 2(d_{I^\dagger \pi(I^\dagger)} + d_{J^\dagger \pi(J^\dagger)} + d_{K^\dagger \pi(K^\dagger)} + d_{L^\dagger \pi(L^\dagger)}) \tag{104}$$

$$T^\dagger\mathbf{1}_5 = 2(d_{I^\dagger K^\dagger} + d_{J^\dagger L^\dagger} + d_{\pi(I^\dagger)\pi(K^\dagger)} + d_{\pi(J^\dagger)\pi(L^\dagger)}). \tag{105}$$

Since we are on $\mathbf{1}_5$, we need to consider $p_3(\cdot)$ as introduced in (63). On $\mathbf{1}_5$, $p_3(\cdot)$ is given by

$$p_3(i, j, k, l, s, t, r) = \frac{1}{(n-1)(n-3)(n-5)} p(i, j, k, l) \quad \text{when } |\{i, j, k, l, s, t, r\}| = 7. \tag{106}$$

The justification for (106) is essentially the same as that for (64).

Using $p_3(\cdot)$, we begin to bound the summands in (104).

$$\begin{aligned}
E|d_{I^\dagger \pi(I^\dagger)}|\mathbf{1}_5 &= \sum_{|\{i,j,k,l,s,t,r\}|=7} |d_{is}| p_3(i, j, k, l, s, t, r) \\
&= \frac{(n-6)(n-5)}{(n-1)(n-3)(n-5)} \sum_{|\{i,j,k,l,s\}|=5} |d_{is}| p(i, j, k, l) \\
&\leq 8 \frac{\beta_D}{n} \quad \text{when } n \geq 9.
\end{aligned} \tag{107}$$

It is easy to see that $I^\dagger, J^\dagger, K^\dagger, L^\dagger$ have identical marginal distributions and since π is chosen independently of these indices, we have $E|d_{N\pi(N)}|\mathbf{1}_5$ is constant over $\{I^\dagger, J^\dagger, K^\dagger, L^\dagger\}$. Thus we obtain

$$E|T\mathbf{1}_5| \leq 64 \frac{\beta_D}{n} \quad \text{when } n \geq 9. \tag{108}$$

Now bounding the L^1 norm of the first summand in (105),

$$\begin{aligned}
E|d_{I^\dagger K^\dagger}|\mathbf{1}_5 &= \sum_{|\{i,j,k,l,s,t,r\}|=7} |d_{ik}| p_3(i, j, k, l, s, t, r) \\
&= \frac{(n-4)(n-5)(n-6)}{(n-1)(n-3)(n-5)} \sum_{|\{i,j,k,l\}|=4} |d_{ik}| p(i, j, k, l) \\
&\leq 4 \frac{\beta_D}{n} \quad \text{when } n \geq 9, \text{ using (43)}.
\end{aligned} \tag{109}$$

Now consider the last summand of (105),

$$\begin{aligned}
E|d_{\pi(J^\dagger)\pi(L^\dagger)}|\mathbf{1}_5 &= \sum_{|\{i,j,k,l,s,t,r\}|=7} |d_{tr}|p_3(i,j,k,l,s,t,r) \\
&\leq \frac{n-6}{(n-1)(n-3)(n-5)} \sum_{|\{i,j,k,l,t,r\}|=6} |d_{tr}|p(i,j,k,l) \\
&\leq 8\frac{\beta_D}{n} \quad \text{when } n \geq 9, \text{ using (45).}
\end{aligned} \tag{110}$$

Since $(I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger), \pi(K^\dagger), \pi(L^\dagger)) \stackrel{C}{=} (J^\dagger, I^\dagger, L^\dagger, K^\dagger, \pi(J^\dagger), \pi(I^\dagger), \pi(L^\dagger), \pi(K^\dagger))$, we see that

$$E|d_{J^\dagger K^\dagger}|\mathbf{1}_5 = E|d_{J^\dagger L^\dagger}|\mathbf{1}_5 \quad \text{and} \quad E|d_{\pi(I^\dagger)\pi(K^\dagger)}|\mathbf{1}_5 = E|d_{\pi(J^\dagger)\pi(L^\dagger)}|\mathbf{1}_5. \tag{111}$$

So, using (109),(110) along with (111), we obtain

$$E|T^\dagger|\mathbf{1}_5 \leq 48\frac{\beta_D}{n} \quad \text{when } n \geq 9. \tag{112}$$

Combining (108) and (112), we obtain

$$EV\mathbf{1}_5 \leq 112\frac{\beta_D}{n} \quad \text{when } n \geq 9. \tag{113}$$

Combining the bounds from (78),(91),(98),(103) and (113), we obtain

$$E|W - W^*| \leq EV \leq 112\frac{\beta_D}{n} + 672\frac{\beta_D}{n^2} + 192\frac{\beta_D}{n^3} \quad \text{when } n \geq 9. \tag{114}$$

This completes the proof of Theorem 4.2. $\square$

5 L^∞ bounds

In this section we will use Theorem 4.2 obtained in the previous section to obtain L^∞ bounds using arguments similar to those in [1]. It is worth noting that we can use L^1 along with L^∞ bounds to obtain L^p bounds for any $p \geq 1$ using (3). The main theorem of this section is the following

Theorem 5.1. *Suppose we have an $n \times n$ array $D = ((d_{ij}))$ satisfying $d_{ij} = d_{ji}, d_{ii} = d_{i+} = 0 \forall i, j$ and $\sigma_D^2 = 1$. If $W = \sum d_{i\pi(i)}$ where π is an involution chosen uniformly at random from Π_n , then for $n > 9$*

$$\|F_W - \Phi\|_\infty \leq K\frac{\beta_D}{n} \tag{115}$$

Here $K = 61,702,446$ is a universal constant.

Theorem 5.1 readily implies Theorem 2.1 by (3) and Theorem 4.1.

We claim that to prove Theorem 5.1 it is enough to consider arrays with $\beta_D/n \leq \epsilon_0 = 1/90$, and $n \geq n_0 = 1000$. To prove the claim, first note that

$$\beta_D^{\frac{1}{3}} = \left(\sum_{1 \leq i,j \leq n} |d_{ij}|^3 \right)^{\frac{1}{3}} \geq n^{-1/3} \left(\sum |d_{ij}|^2 \right)^{\frac{1}{2}} = n^{-1/3} \left(\frac{(n-1)(n-3)}{2(n-2)} \right)^{\frac{1}{2}}, \tag{116}$$

which is greater than $1/2$ for $n \geq 4$. In (116), the first inequality follows from Hölder's inequality and the next equality follows from the fact that $\sigma_D^2 = 1$. So if $n < n_0$ then $\beta_D/n > 1/(8n_0)$. Since $K > \max\{1/\epsilon_0, 8n_0\}$ inequality (115) holds if $\beta_D/n > \epsilon_0$ or $n < n_0$.

One useful inequality that will be used repeatedly in the proof is the following

$$\left(\left| \sum_{i=1}^k a_i \right| \right)^3 \leq \left(\sum |a_i| \right)^3 \leq k^2 \sum |a_i|^3. \quad (117)$$

The proof of Theorem 5.1 proceeds by first proving several auxiliary lemmas. The first lemma helps bound the error created when truncating the array as in [1], page 382. For the array $D = ((d_{ij}))_{n \times n}$, we define

$$d'_{ij} = d_{ij} 1_{|d_{ij}| \leq \frac{1}{2}}. \quad (118)$$

Letting $\Gamma = \{(i, j) : |d_{ij}| > \frac{1}{2}\}$ and $\Gamma_i = \{j : (i, j) \in \Gamma\}$ we have

$$|\Gamma_i| \leq 8 \sum_j |d_{ij}|^3 \quad \text{and hence} \quad (119)$$

$$|\Gamma| \leq 8\beta_D. \quad (120)$$

Inequality (120) has the following useful consequence, that is,

$$\begin{aligned} P(Y_{D'} \neq Y_D) &\leq P\left(\sum_i 1_{\Gamma}(i, \pi(i)) \geq 1\right) \\ &\leq E\left(\sum_i 1_{\Gamma}(i, \pi(i))\right) \\ &= \frac{|\Gamma|}{n-1} < \frac{2|\Gamma|}{n} \leq \frac{16\beta_D}{n}. \end{aligned} \quad (121)$$

Lemma 5.1. *Suppose $((d_{ij}))_{n \times n}$ satisfies the conditions in Theorem 5.1. Then with $((d'_{ij}))_{n \times n}$ defined in (118), $\beta_D \leq \epsilon_0 n$ and $n \geq n_0$, we have,*

$$|d'_{++}| \leq 4\beta_D \quad \text{and} \quad |d'_{i+}| \leq 4 \sum_j |d_{ij}|^3 \leq 4\beta_D, \quad (122)$$

and therefore

$$\begin{aligned} |\mu_{D'}| &\leq 8 \frac{\beta_D}{n}, \\ |\sigma_{D'}^2 - 1| &\leq 10 \frac{\beta_D}{n} \quad \text{and} \quad \beta_{D'} \leq 22\beta_D. \end{aligned}$$

Proof. Using $d_{++} = 0$ and (120), we have

$$|d'_{++}| = \left| \sum_{(i,j): |d_{ij}| \leq 1/2} d_{ij} \right| = \left| \sum_{(i,j) \in \Gamma} d_{ij} \right| \leq \sum_{(i,j) \in \Gamma} |d_{ij}| \leq |\Gamma|^{\frac{2}{3}} \beta_D^{\frac{1}{3}} \leq 4\beta_D. \quad (123)$$

Similarly, as $d_{i+} = 0$ for all $i \in \{1, \dots, n\}$, we obtain using (119),

$$|d'_{i+}| = \left| \sum_{j \notin \Gamma_i} d_{ij} \right| = \left| \sum_{j \in \Gamma_i} d_{ij} \right| \leq |\Gamma_i|^{2/3} \left(\sum_{j \in \Gamma_i} |d_{ij}|^3 \right)^{1/3} \leq 4 \sum_j |d_{ij}|^3 \leq 4\beta_D. \quad (124)$$

Now, using (5) and (123),

$$|\mu_{D'}| = \left| \frac{d'_{++}}{n-1} \right| \leq \frac{4\beta_D}{n-1} \leq 8 \frac{\beta_D}{n}.$$

To prove the last assertion, first note that by (124) we have

$$\sum_{i=1}^n |d'_{i+}|^2 \leq 4\beta_D \sum_{i=1}^n |d'_{i+}| \leq 16\beta_D \sum_i \sum_j |d_{ij}|^3 = 16\beta_D^2. \quad (125)$$

Similarly one can obtain

$$\sum_{i=1}^n |d'_{i+}|^3 \leq 64\beta_D^3. \quad (126)$$

From (5) and the fact that $\sigma_D^2 = 1$ we obtain the following,

$$\begin{aligned} |\sigma_{D'}^2 - 1| &= \left| \frac{2}{(n-1)(n-3)} \left((n-2) \sum_{1 \leq i,j \leq n} (d_{ij}^2 - d'_{ij}^2) + \frac{1}{n-1} d_{++}^2 - 2 \sum_{i=1}^n d_{i+}^2 \right) \right| \\ &\leq \frac{2}{(n-1)(n-3)} \left((n-2) \sum_{(i,j) \in \Gamma} d_{ij}^2 + \frac{1}{n-1} d_{++}^2 + 2 \sum_{i=1}^n d_{i+}^2 \right) \\ &\leq \frac{2}{(n-1)(n-3)} \left((n-2) \sum_{(i,j) \in \Gamma} d_{ij}^2 + 16 \frac{\beta_D^2}{(n-1)} + 32\beta_D^2 \right) \quad \text{using (123) and (125)} \quad (127) \\ &\leq 8 \frac{\beta_D}{n-1} + 32 \frac{\beta_D^2}{(n-1)^2(n-3)} + 64 \frac{\beta_D^2}{(n-1)(n-3)} \quad \text{since } \sum_{(i,j) \in \Gamma} d_{ij}^2 \leq \beta_D^{\frac{2}{3}} |\Gamma|^{\frac{1}{3}} \leq 2\beta_D \\ &\leq 10 \frac{\beta_D}{n} < \frac{1}{9} \quad \text{since } \beta_D/n \leq 1/90 \text{ and } n \geq 1000. \quad (128) \end{aligned}$$

Hence, we obtain $|\sigma_{D'}^2 - 1| < 1/9$. Thus using (6) with notation as in (11), and inequality (117), we obtain

$$\begin{aligned} \beta_{D'} &= \frac{1}{\sigma_{D'}^3} \sum_{i \neq j} \left| d'_{ij} - \frac{d'_{i+}}{n-2} - \frac{d'_{+j}}{n-2} + \frac{d'_{++}}{(n-1)(n-2)} \right|^3 \\ &\leq 16 \frac{1}{\sigma_{D'}^3} \sum_{i \neq j} \left(|d'_{ij}|^3 + \frac{|d'_{i+}|^3}{(n-2)^3} + \frac{|d'_{+j}|^3}{(n-2)^3} + \frac{|d'_{++}|^3}{((n-1)(n-2))^3} \right) \\ &\leq 16 \frac{1}{\sigma_{D'}^3} \left(\beta_D + 128 \frac{\beta_D^3}{(n-2)^3} + 64n^2 \frac{\beta_D^3}{((n-1)(n-2))^3} \right) \\ &\leq 22\beta_D \quad \text{since } |\sigma_{D'}^2 - 1| < 1/9, \beta_D/n \leq 1/90, \end{aligned}$$

where in the second inequality we use (123) and (126). $\square$

Lemma 5.2. Let $D = ((d_{ij}))$ be an array as in Theorem 5.1 and $D' = ((d'_{ij}))$ be as in (118). Let us denote

$$\tilde{d}_{ij} = \frac{\widehat{d'_{ij}}}{\sigma_{D'}}.$$

If $\beta_D/n \leq \epsilon_0$ and $n \geq n_0$, we have $|\tilde{d}_{ij}| \leq 1$.

Proof. Because $\sigma_{D'}^2 > 1/2$ from Lemma 5.1, $\sigma_{D'} > 2/3$. Using this inequality, definition (6) and (122) we

obtain

$$\begin{aligned}
|\tilde{d}_{ij}| &\leq \frac{1}{\sigma_{D'}} \left(|d'_{ij}| + \frac{1}{n-2} (|d'_{i+}| + |d'_{+j}|) + \frac{|d'_{++}|}{(n-1)(n-2)} \right) \\
&\leq \frac{3}{4} + \frac{8}{\sigma_{D'}} \frac{\beta_D}{n-2} + \frac{4}{\sigma_{D'}} \frac{\beta_D}{(n-1)(n-2)} \\
&< \frac{3}{4} + 16 \frac{\beta_D}{n} + 4 \frac{\beta_D}{n} \quad \text{since } n \geq 1000 \\
&= \frac{3}{4} + 20 \frac{\beta_D}{n} < 1 \quad \text{since } \beta_D/n \leq 1/90.
\end{aligned}$$

This completes the proof. $\square$

For $E = ((e_{ij}))_{n \times n}$ let F_E be the distribution function of Y_E and

$$\delta_E = \|F_E - \Phi\|_\infty. \quad (129)$$

For $\gamma > 0$ let $M_n(\gamma)$ be the set of $n \times n$ matrices $E = ((e_{ij}))$ satisfying

$$\sigma_E^2 = 1, e_{ij} = e_{ji}, e_{ii} = e_{i+} = 0 \quad \forall i, j \quad \text{and} \quad \beta_E \leq \gamma.$$

Let us define

$$\delta(\gamma, n) = \sup_{E \in M_n(\gamma)} \delta_E. \quad (130)$$

Also, we define

$$\delta^1(\gamma, n) = \sup_{E \in M_n^1(\gamma)} \delta_E \quad \text{where} \quad M_n^1(\gamma) = \{E \in M_n(\gamma) : \sup_{i,j} |e_{ij}| \leq 1\}. \quad (131)$$

Lemma 5.3. *When $n \geq n_0$, with δ_D and $\delta^1(\gamma, n)$ defined in (129) and (131), for all $\gamma > 0$*

$$\sup\{\delta_D : D \in M_n(\gamma), \beta_D/n \leq \epsilon_0\} \leq \delta^1(22\gamma, n) + 36 \frac{\gamma}{n}.$$

Proof. Let $D = ((d_{ij})) \in M_n(\gamma)$ with $\beta_D/n \leq 1/90$. Hence, by (121) and Lemmas 5.2 and 5.1,

$$\begin{aligned}
\delta_D = \|F_D - \Phi\|_\infty &\leq \sup_t |P(Y_{D'} \leq t) - \Phi(t)| + P(Y_D \neq Y_{D'}) \\
&\leq \sup_t |P(Y_{D'} \leq t) - \Phi(t)| + \frac{16\beta_D}{n} \\
&\leq \delta^1(22\beta_D, n) + \sup_t \left| \Phi\left(\frac{t - \mu_{D'}}{\sigma_{D'}}\right) - \Phi(t) \right| + \frac{16\beta_D}{n}.
\end{aligned} \quad (132)$$

Since $\delta^1(\gamma, n)$ is monotone in γ ,

$$\delta^1(22\beta_D, n) \leq \delta^1(22\gamma, n) \quad \text{when } \beta_D \leq \gamma;$$

similarly, we bound the last term as $16\beta_D/n \leq 16\gamma/n$. So, we are only left with verifying that the second term is bounded by $20\gamma/n$.

From Lemma 5.1, we obtain $|\sigma_{D'}^2 - 1| \leq 1/9$ yielding in particular $\sigma_{D'} \in [2/3, 4/3]$. First consider the case where $|t| \geq 8\beta_D/n$, and for a given t let $t_1 = (t - \mu_{D'})/\sigma_{D'}$. Since $|\mu_{D'}| \leq 8\beta_D/n$ by Lemma 5.1, t and t_1 will be on the same side of the origin. Next, it is easy to show that for $a > 0$ we have $|t \exp(-at^2/2)| \leq 1/\sqrt{a}$. Hence

$$\begin{aligned}
\left| t \exp\left(-\frac{9}{32}(t - \mu_{D'})^2\right) \right| &\leq \left| (t - \mu_{D'}) \exp\left(-\frac{9}{32}(t - \mu_{D'})^2\right) \right| + |\mu_{D'}| \\
&\leq \frac{4}{3} + |\mu_{D'}| \\
&\leq \frac{4}{3}(1 + |\mu_{D'}|).
\end{aligned} \quad (133)$$

Since $\sigma_{D'} \geq 2/3$ and Lemma 5.1 gives $|\sigma_{D'}^2 - 1| \leq 10\beta_D/n$, we find that

$$|\sigma_{D'} - 1| = \frac{|\sigma_{D'}^2 - 1|}{\sigma_{D'} + 1} \leq 10 \frac{\beta_D}{n}.$$

Now, by the mean value theorem, $\sigma_{D'} \in [2/3, 4/3]$, and Lemma 5.1

$$\begin{aligned} |\Phi(t_1) - \Phi(t)| &\leq \max \left(\phi\left(\frac{t - \mu_{D'}}{\sigma_{D'}}\right), \phi(t) \right) \left| \frac{t - \mu_{D'}}{\sigma_{D'}} - t \right| \quad \text{where } \phi = \Phi' \\ &\leq \frac{1}{\sqrt{2\pi}} \max \left\{ \exp\left(-\frac{9}{32}(t - \mu_{D'})^2\right), \exp\left(-\frac{t^2}{2}\right) \right\} \left| \frac{t(1 - \sigma_{D'})}{\sigma_{D'}} \right| + \frac{1}{\sqrt{2\pi}} \left| \frac{\mu_{D'}}{\sigma_{D'}} \right| \\ &\leq \frac{3}{2\sqrt{2\pi}} |\sigma_{D'} - 1| \max \left\{ \left| t \exp\left(-\frac{9}{32}(t - \mu_{D'})^2\right) \right|, \left| t \exp\left(-\frac{t^2}{2}\right) \right| \right\} + \frac{1}{\sqrt{2\pi}} \left| \frac{\mu_{D'}}{\sigma_{D'}} \right| \\ &\leq \frac{3}{2\sqrt{2\pi}} |\sigma_{D'} - 1| \left(\frac{4}{3}(1 + |\mu_{D'}|) \right) + \frac{3}{4} |\mu_{D'}| \quad \text{using (133)} \\ &\leq \frac{2}{\sqrt{2\pi}} |\sigma_{D'} - 1| (1 + |\mu_{D'}|) + \frac{3}{4} |\mu_{D'}| \\ &\leq \frac{2}{\sqrt{2\pi}} 10 \frac{\beta_D}{n} (1 + 8 \frac{\beta_D}{n}) + 6 \frac{\beta_D}{n} \\ &\leq 17 \frac{\beta_D}{n} \quad \text{since } \beta_D/n \leq 1/90. \end{aligned} \tag{134}$$

When $|t| < 8\beta_D/n$, the bound is easier. Since t_1 lies in the interval with boundary points $3(t - \mu_{D'})/2$ and $3(t - \mu_{D'})/4$, we have

$$|t_1| \leq \frac{3(|t| + |\mu_{D'}|)}{2}. \tag{135}$$

Now, using $|t| < 8\beta_D/n$, $|\mu_{D'}| < 8\beta_D/n$ and (135), we obtain

$$\begin{aligned} |\Phi(t_1) - \Phi(t)| &\leq \frac{1}{\sqrt{2\pi}} |t_1 - t| \\ &\leq \frac{1}{\sqrt{2\pi}} (3|t| + 2|\mu_{D'}|) \\ &\leq \frac{1}{\sqrt{2\pi}} 40 \frac{\beta_D}{n} < 20 \frac{\beta_D}{n}. \end{aligned}$$

Thus we obtain

$$\sup_t \left| \Phi\left(\frac{t - \mu_{D'}}{\sigma_{D'}}\right) - \Phi(t) \right| \leq 20 \frac{\beta_D}{n} \leq 20 \frac{\gamma}{n},$$

completing the proof. $\square$

In view of Lemma 5.3, to prove Theorem 5.1 it remains only to show $\delta^1(\gamma, n) \leq c\gamma/n$ for an explicit c which we eventually determine. Hence in the following calculations we consider only arrays $D = ((d_{ij}))_{n \times n}$ in $M_n^1(\gamma)$, and so $\sup_{i,j} |d_{ij}| \leq 1$.

We will need the following technical lemma.

Lemma 5.4. *If $D \in M_n^1(\gamma)$ and $B = ((b_{ij}))_{n-l \times n-l}$ is the array formed by removing from D the l rows and l columns indexed by $\mathcal{T} \subset \{1, 2, \dots, n\}$, where l is even and $l = |\mathcal{T}| \leq 8$, then for $n \geq n_0$ and $\beta_D/n \leq \epsilon_0/50$, $|\mu_B| \leq 8.07$, $|\sigma_B^2 - 1| \leq .52$, and $\beta_B \leq 50\beta_D$.*

Proof. To prove the first claim, we note since $d_{i+} = 0$, $|d_{ij}| \leq 1$ and $l \leq 8$, letting $\bar{n} = n - l$,

$$|b_{i+}| = \left| \sum_{j \notin \mathcal{T}} d_{ij} \right| = \left| - \sum_{j \in \mathcal{T}} d_{ij} \right| \leq 8 \quad \text{and hence} \quad |b_{++}| = \left| \sum_{i \notin \mathcal{T}} b_{i+} \right| \leq 8\bar{n}. \quad (136)$$

This inequality implies $|\mu_B| = |b_{++}|/(\bar{n} - 1) \leq 8.07$ proving the first claim.

Using $\sigma_D^2 = 1$ and (5) we obtain

$$2 \frac{\bar{n} - 2}{(\bar{n} - 1)(\bar{n} - 3)} \sum_{1 \leq i, j \leq n} d_{ij}^2 = \frac{(\bar{n} - 2)(n - 1)(n - 3)}{(n - 2)(\bar{n} - 1)(\bar{n} - 3)} \in [1, 1.01] \quad \text{when } n \geq 1000. \quad (137)$$

Using the above bound, (5) and (136), we obtain

$$\left| \sigma_B^2 - 2 \frac{\bar{n} - 2}{(\bar{n} - 1)(\bar{n} - 3)} \sum_{1 \leq i, j \leq n} d_{ij}^2 \right| \quad (138)$$

$$\begin{aligned} &\leq \frac{2}{(\bar{n} - 1)(\bar{n} - 3)} \left((\bar{n} - 2) \left| \sum_{\{i \notin \mathcal{T}\} \cap \{j \notin \mathcal{T}\}} d_{ij}^2 - \sum_{i, j=1}^n d_{ij}^2 \right| + \frac{1}{\bar{n} - 1} b_{++}^2 + 2 \sum_{i \notin \mathcal{T}} b_{i+}^2 \right) \\ &\leq \frac{2}{(\bar{n} - 1)(\bar{n} - 3)} \left((\bar{n} - 2) \sum_{\{i \in \mathcal{T}\} \cup \{j \in \mathcal{T}\}} d_{ij}^2 + \frac{64\bar{n}^2}{\bar{n} - 1} + 128\bar{n} \right) \\ &\leq \frac{32n}{272(\bar{n} - 3)} + \frac{128\bar{n}^2}{(\bar{n} - 1)^2(n - 3)} + \frac{256\bar{n}}{(\bar{n} - 1)(\bar{n} - 3)} \\ &\leq .51 \quad \text{when } n \geq 1000, \end{aligned} \quad (139)$$

$$\leq .51 \quad \text{when } n \geq 1000, \quad (140)$$

where for (139), we use $|\mathcal{T}| \leq 8$, $\beta_D \leq n/(50 \times 90)$ and Hölder's inequality to obtain

$$\begin{aligned} \sum_{\{i \in \mathcal{T}\} \cup \{j \in \mathcal{T}\}} d_{ij}^2 &\leq \sum_{i \in \mathcal{T}} \sum_{j=1}^n d_{ij}^2 + \sum_{j \in \mathcal{T}} \sum_{i=1}^n d_{ij}^2 = 2 \sum_{i \in \mathcal{T}} \sum_{j=1}^n d_{ij}^2 \\ &\leq 2 \times 8 \beta_D^{\frac{2}{3}} n^{\frac{1}{3}} \quad \text{since } \sum_{j=1}^n d_{ij}^2 \leq (\sum_{j=1}^n |d_{ij}|^3)^{\frac{2}{3}} n^{\frac{1}{3}} \leq \beta_D^{\frac{2}{3}} n^{\frac{1}{3}} \text{ and } |\mathcal{T}| \leq 8 \\ &\leq 16 \frac{n}{272} \quad \text{since } \beta_D \leq n/4500. \end{aligned}$$

From (140) and (137), we obtain $|\sigma_B^2 - 1| \leq .52$ for $n \geq 1000$, proving the second claim of the lemma.

To prove the final claim we observe $|b_{i+}| = |\sum_{j \in \mathcal{T}} d_{ij}|$. Thus, by (117) and $|\mathcal{T}| \leq 8$,

$$\begin{aligned} \sum_{i \notin \mathcal{T}} |b_{i+}|^3 &= \sum_{i \notin \mathcal{T}} \left| \sum_{j \in \mathcal{T}} d_{ij} \right|^3 \leq 64 \sum_{i \notin \mathcal{T}} \sum_{j \in \mathcal{T}} |d_{ij}|^3 \leq 64\beta_D \quad \text{and similarly} \\ |b_{++}|^3 &= \left| \sum_{i \notin \mathcal{T}} b_{i+} \right|^3 \leq \bar{n}^2 \sum_{i \notin \mathcal{T}} |b_{i+}|^3 \leq 64\bar{n}^2 \beta_D. \end{aligned} \quad (141)$$

These observations, along with the fact that $\sigma_B^2 \geq .48$, yield

$$\begin{aligned} \beta_B &= \frac{\sum_{\{i \notin \mathcal{T}\} \cap \{j \notin \mathcal{T}\}} |\hat{b}_{ij}|^3}{\sigma_B^3} \leq 3.1 \sum_{\{i \notin \mathcal{T}\} \cap \{j \notin \mathcal{T}\}} |\hat{b}_{ij}|^3 \\ &= 3.1 \sum_{\{i \notin \mathcal{T}\} \cap \{j \notin \mathcal{T}\}} \left| b_{ij} - \frac{b_{i+}}{(\bar{n} - 2)} - \frac{b_{+j}}{(\bar{n} - 2)} + \frac{b_{++}}{(\bar{n} - 1)(\bar{n} - 2)} \right|^3 \\ &\leq 3.1 \times 4^2 \left(\sum_{i \notin \mathcal{T}} |b_{ij}|^3 + 2\bar{n} \sum_{i \notin \mathcal{T}} \frac{|b_{i+}|^3}{(\bar{n} - 2)^3} + \bar{n}^2 \frac{|b_{++}|^3}{((\bar{n} - 1)(\bar{n} - 2))^3} \right) \quad \text{using (117)} \\ &\leq 49.6 \left(\beta_D + 128 \frac{\bar{n}}{(\bar{n} - 2)^3} \beta_D + 64\beta_D \frac{\bar{n}^4}{((\bar{n} - 1)(\bar{n} - 2))^3} \right) \quad \text{using (141)} \\ &\leq 50\beta_D \quad \text{when } n \geq 1000, \text{ since } l \leq 8. \end{aligned}$$

This completes the proof. $\square$

Lemma 5.5. Consider a nonnegative sequence $\{a_n, n \in \mathbb{Z}\}$ such that $a_n = 0$ for all $n \leq 0$ and

$$a_n \leq c + \alpha \max_{l \in \{4,6,8\}} a_{n-l} \quad \text{for all } n \geq 1, \quad (142)$$

where $c \geq 0$ and $\alpha \in (0, 1/3)$. Then, with $b = \max\{c, a_1(1 - 3\alpha)\}$, for all $n \geq 1$

$$a_n \leq \frac{b}{1 - 3\alpha}.$$

Proof. Letting $b_n, n \geq 1$ be given by the recursion

$$b_{n+1} = 3\alpha b_n + b \quad \text{for } n \geq 1, \text{ with } b_1 = a_1,$$

explicitly solving yields for $n > 1$

$$b_n = c_1(3\alpha)^n + c_2 \quad \text{where } c_1 = \frac{a_1(1-3\alpha)-b}{3\alpha(1-3\alpha)} \text{ and } c_2 = \frac{b}{1-3\alpha}.$$

Since $c_1 \leq 0$ and $3\alpha < 1$, b_n is increasing and bounded above by c_2 . Hence it suffices to show $a_n \leq b_n$.

Since a_n is nonnegative, (142) implies that

$$a_n \leq c + \alpha \sum_{l \in \{4,6,8\}} a_{n-l} \quad \text{for all } n \geq 1. \quad (143)$$

We show $a_m \leq b_m$ for all $1 \leq m \leq n$ by induction. When $n \leq 4$, since $a_{n-l} = 0$ for $l \in \{4,6,8\}$ we have

$$a_n \leq c \leq b \leq b_n.$$

Now supposing the claim is true for some $n \geq 4$, by (143) and the monotonicity of $b_n, n \geq 1$, we obtain

$$a_{n+1} \leq c + \alpha \sum_{l \in \{4,6,8\}} a_{n+1-l} \leq b + \alpha \sum_{l \in \{4,6,8\}} b_{n+1-l} \leq b + 3\alpha b_{n-3} \leq b + 3\alpha b_n = b_{n+1}.$$

This completes the proof. $\square$

Lemma 5.6. With $\delta^1(\gamma, n)$ defined as in (131),

$$\delta^1(\gamma, n) \leq 2804655 \frac{\gamma}{n} \quad \text{when } n \geq n_0 = 1000.$$

Proof. We will consider a smoothed family of indicator functions indexed by $\lambda > 0$, namely

$$h_{z,\lambda}(x) = \begin{cases} 1 & \text{if } x \leq z \\ 1 + (z - x)/\lambda & \text{if } x \in (z, z + \lambda] \\ 0 & \text{if } x > z + \lambda. \end{cases} \quad (144)$$

Also, define

$$h_{z,0}(x) = 1_{(-\infty, z]}(x).$$

Let $f_{z,\lambda}$ denote the solution to the following Stein equation

$$f'(x) - xf(x) = h_{z,\lambda} - \Phi(h_{z,\lambda}) \quad \text{where } \Phi(h_{z,\lambda}) = E(h_{z,\lambda}(Z)) \text{ and } Z \sim \mathcal{N}(0, 1). \quad (145)$$

We will need the following key inequality about $f_{z,\lambda}$ from [1],

$$|f'_{z,\lambda}(x+y) - f'_{z,\lambda}(x)| \leq |y| \left(1 + 2|x| + \frac{1}{\lambda} \int_0^1 1_{[z, z+\lambda]}(x+ry) dr \right) \quad \text{for any } \lambda > 0. \quad (146)$$

We consider $D \in M_n^1(\gamma)$ with $\beta_D/n \leq \epsilon_0/51$ and $n \geq 1000$. Let $W = \sum_{i=1}^n d_{i\pi(i)}$ as before. Using the fact that

$$h_{z,0} \leq h_{z,\lambda} \leq h_{z+\lambda,0},$$

and recalling the definition of δ_D in (129), we obtain

$$\delta_D \leq \sup_z |E(h_{z,\lambda}(W)) - \Phi(h_{z,\lambda})| + \lambda/\sqrt{2\pi}. \quad (147)$$

From (145), (12) and $\text{Var}(W) = 1$, we obtain

$$\begin{aligned} \sup_z |E(h_{z,\lambda}(W)) - \Phi(h_{z,\lambda})| &= \sup_z |E(f'_{z,\lambda}(W)) - E(Wf_{z,\lambda}(W))| \\ &= \sup_z |E(f'_{z,\lambda}(W)) - E(f'_{z,\lambda}(W^*))| \\ &\leq \sup_z E|f'_{z,\lambda}(W^*) - f'_{z,\lambda}(W)|. \end{aligned} \quad (148)$$

From (147), (148) and (146), we obtain the following

$$\begin{aligned} \delta_D &\leq \sup_z E|f'_{z,\lambda}(W^*) - f'_{z,\lambda}(W)| + \lambda/\sqrt{2\pi} \\ &\leq \sup_z E \left\{ |W^* - W| \left(1 + 2|W| + \frac{1}{\lambda} \int_0^1 1_{[z, z+\lambda]}(W + r(W^* - W)) dr \right) \right\} + \lambda/\sqrt{2\pi} \\ &= E|W^* - W| + 2E(|W||W^* - W|) \\ &\quad + \sup_z \frac{1}{\lambda} E \left(|W^* - W| \int_0^1 1_{[z, z+\lambda]}(W + r(W^* - W)) dr \right) + \lambda/\sqrt{2\pi} \\ &:= A_1 + A_2 + A_3 + \lambda/\sqrt{2\pi} \quad \text{say.} \end{aligned} \quad (149)$$

First, A_1 is bounded using Theorem 4.2, as follows

$$\begin{aligned} A_1 &= E|W^* - W| \leq 112 \frac{\beta_D}{n} + 672 \frac{\beta_D}{n^2} + 192 \frac{\beta_D}{n^3} \\ &\leq 113 \frac{\beta_D}{n} \quad \text{since } n \geq n_0 = 1000. \end{aligned} \quad (150)$$

Next we bound A_2 . From Theorem 4.2, we obtain,

$$\begin{aligned} W^* - W &= (UW^\dagger + (1 - U)W^\ddagger) - W \\ &= (U(S + T^\dagger) + (1 - U)(S + T^\ddagger)) - (S + T) = UT^\dagger + (1 - U)T^\ddagger - T. \end{aligned} \quad (151)$$

Let $\mathbf{I} = (I^\dagger, J^\dagger, K^\dagger, L^\dagger, \pi(I^\dagger), \pi(J^\dagger), \pi(K^\dagger), \pi(L^\dagger))$ and $\mathcal{I}$ be as defined in Lemma 3.2. Thus, by (47), the right hand side of (151), and hence $W^* - W$, is measurable with respect to $\mathcal{J} = \{\mathbf{I}, U\}$. Furthermore, since $\sup_{i,j} |d_{ij}| \leq 1$ and $|Z| \leq 8$, we have

$$|W| = |S + T| \leq |S| + |T| = |S| + \left| \sum_{i \in \mathcal{I}} d_{i\pi(i)} \right| \leq |S| + \sum_{i \in \mathcal{I}} |d_{i\pi(i)}| \leq |S| + 8.$$

Now, by the definition of A_2 , and that U is independent of S and $\mathbf{I}$,

$$\begin{aligned} A_2 &= 2E(|W||W^* - W|) \\ &= 2E(|W^* - W|E(|W||\mathcal{J})) \leq 2E(|W^* - W|E(|S| + 8|\mathcal{J}|)) \\ &\leq 2E(|W^* - W|\sqrt{E(S^2|\mathbf{I})}) + 16E|W^* - W|. \end{aligned} \quad (152)$$

In the following, for $\mathbf{i}$ a realization of $\mathbf{I}$, let $|\mathbf{i}|$ denote the number of distinct elements of $\mathbf{i}$. Since $S = \sum_{i \notin \mathcal{I}} d_{i\pi(i)}$ and π is chosen from Π_n uniformly at random, conditioned on $\mathbf{I} = \mathbf{i}$, S has the same distribution

as $\sum_{i \notin \mathbf{i}} b_{i\theta(i)}$, where $B = ((b_{ij}))_{n-|\mathbf{i}| \times n-|\mathbf{i}|}$ is the matrix formed by removing rows and columns of D that occur in $\mathbf{i}$ and θ is chosen uniformly from $\Pi_{n-|\mathbf{i}|}$. Since $|\mathbf{i}| \leq 8$ and $\beta_D/n \leq \epsilon_0/51 < \epsilon_0/50$, Lemma 5.4 yields $|\mu_B| \leq 8.07$ and $\sigma_B^2 \leq 1.52$, and hence

$$E|Y_B|^2 \leq 1.52 + 8.07^2 = 66.6449 \quad \text{when } n \geq 1000. \quad (153)$$

Using (153), we obtain

$$E(S^2 | \mathbf{I} = \mathbf{i}) = E|Y_B|^2 \leq 67 \quad \text{for all } \mathbf{i}.$$

Thus using (152) and (150), we obtain

$$A_2 \leq 33E|W^* - W| \leq 3729 \frac{\beta_D}{n}. \quad (154)$$

Finally, we are left with bounding A_3 . First we note that

$$\begin{aligned} W + r(W^* - W) &= rW^* + (1-r)W \\ &= r(S + UT^\dagger + (1-U)T^\ddagger) + (1-r)(S + T) \\ &= S + rUT^\dagger + r(1-U)T^\ddagger + (1-r)T \\ &= S + g_r \quad \text{where } g_r = rUT^\dagger + r(1-U)T^\ddagger + (1-r)T. \end{aligned}$$

Now, from the definition of A_3 , again using that $W - W^*$ is $\mathcal{J}$ measurable,

$$\begin{aligned} A_3 &= \sup_z \frac{1}{\lambda} E \left(|W - W^*| \int_0^1 1_{[z, z+\lambda]}(W + r(W^* - W)) dr \right) \\ &= \sup_z \frac{1}{\lambda} E \left(|W - W^*| E \left(\int_0^1 1_{[z, z+\lambda]}(W + r(W^* - W)) dr \middle| \mathcal{J} \right) \right) \\ &= \sup_z \frac{1}{\lambda} E \left(|W - W^*| \int_0^1 P(W + r(W^* - W) \in [z, z + \lambda] | \mathcal{J}) dr \right) \\ &= \sup_z \frac{1}{\lambda} E(|W - W^*| \int_0^1 P(S + g_r \in [z, z + \lambda] | \mathcal{J}) dr) \\ &= \sup_z \frac{1}{\lambda} E(|W - W^*| \int_0^1 P(S \in [z - g_r, z + \lambda - g_r] | \mathcal{J}) dr) \\ &\leq \frac{1}{\lambda} E(|W - W^*| \int_0^1 \sup_z P(S \in [z - g_r, z + \lambda - g_r] | \mathcal{J}) dr) \\ &= \frac{1}{\lambda} E(|W - W^*| \int_0^1 \sup_z P(S \in [z, z + \lambda] | \mathcal{J}) dr) \\ &= \frac{1}{\lambda} E(|W - W^*| \sup_z P(S \in [z, z + \lambda] | \mathcal{J})) \\ &= \frac{1}{\lambda} E(|W - W^*| \sup_z P(S \in [z, z + \lambda] | \mathbf{I})), \end{aligned} \quad (155)$$

where to obtain equality in (155) we have used the fact that g_r is measurable with respect to $\mathcal{J}$ for all r .

It remains only to bound $P(S \in [z, z + \lambda] | \mathbf{I})$. In the following calculations $\tilde{b}_{ij} = \hat{b}_{ij}/\sigma_B$ as before. Since

$\beta_D/n \leq \epsilon_0/51$, Lemma 5.4 yields $\sigma_B > 1/2$. Hence,

$$\begin{aligned}
\sup_z P(S \in [z, z + \lambda] | \mathbf{I} = \mathbf{i}) &= \sup_z P\left(\sum_{i \notin \mathbf{i}} b_{i\theta(i)} \in [z, z + \lambda]\right) \\
&= \sup_z P\left(\sum_{i \notin \mathbf{i}} \frac{b_{i\theta(i)}}{\sigma_B} \in \left[\frac{z}{\sigma_B}, \frac{z + \lambda}{\sigma_B}\right]\right) \\
&\leq \sup_z P\left(\sum_{i \notin \mathbf{i}} \frac{b_{i\theta(i)}}{\sigma_B} \in \left[\frac{z}{\sigma_B}, \frac{z}{\sigma_B} + 2\lambda\right]\right) \\
&= \sup_z P\left(\sum_{i \notin \mathbf{i}} \frac{b_{i\theta(i)}}{\sigma_B} \in [z, z + 2\lambda]\right) \\
&= \sup_z P\left(\sum_{i \notin \mathbf{i}} \tilde{b}_{i\theta(i)} \in [z, z + 2\lambda]\right) \\
&= \sup_z P(Y_{\tilde{B}} \in [z, z + 2\lambda]).
\end{aligned} \tag{157}$$

The equality (157) holds as when computing $\hat{b}_{ij}$ we have that $\sum_{i \notin \mathbf{i}} b_{i+}$ and $\sum_{i \notin \mathbf{i}} b_{i+\theta(i)} = \sum_{j \notin \mathbf{i}} b_{+j}$ do not depend on θ . Recalling that the distribution function of $Y_{\tilde{B}}$ is denoted by $F_{\tilde{B}}$, we have, from the definition of $\delta(\gamma, n)$ in (130),

$$\begin{aligned}
P(Y_{\tilde{B}} \in [z, z + 2\lambda]) &\leq |F_{\tilde{B}}(z + 2\lambda) - \Phi(z + 2\lambda)| + |F_{\tilde{B}}(z) - \Phi(z)| + |\Phi(z + 2\lambda) - \Phi(z)| \\
&\leq 2\delta_{\tilde{B}} + \frac{2\lambda}{\sqrt{2\pi}}.
\end{aligned} \tag{158}$$

Note that $\beta_{\tilde{B}} = \beta_B$ and by Lemma 5.4

$$\frac{\beta_{\tilde{B}}}{n - |\mathbf{i}|} = \frac{\beta_B}{n - |\mathbf{i}|} \leq 50 \frac{\beta_D}{n - |\mathbf{i}|} \leq 51 \frac{\beta_D}{n} \leq \epsilon_0.$$

Hence using the fact that $\tilde{B} \in M_{n-|\mathbf{i}|}(\beta_B)$ and applying Lemma 5.3, we obtain for $n \geq 1008$

$$\delta_{\tilde{B}} \leq \delta^1(22\beta_{\tilde{B}}) + 36 \frac{\beta_{\tilde{B}}}{n - |\mathbf{i}|} = \delta^1(22\beta_B, n - |\mathbf{i}|) + 36 \frac{\beta_B}{n - |\mathbf{i}|}. \tag{159}$$

Inequalities (158) and (159) imply

$$\begin{aligned}
P(Y_{\tilde{B}} \in [z, z + 2\lambda]) &\leq 2\delta_{\tilde{B}} + \frac{2\lambda}{\sqrt{2\pi}} \\
&\leq 2\delta^1(22\beta_B, n - |\mathbf{i}|) + 72 \frac{\beta_B}{n - |\mathbf{i}|} + \frac{2\lambda}{\sqrt{2\pi}} \\
&\leq 2\delta^1(22 \times 50\beta_D, n - |\mathbf{i}|) + \frac{72 \times 50\beta_D}{n - |\mathbf{i}|} + \frac{2\lambda}{\sqrt{2\pi}} \\
&\leq 2 \max_{l \in \{4, 6, 8\}} \delta^1(k_1\beta_D, n - l) + \frac{k_2\beta_D}{n} + \frac{2\lambda}{\sqrt{2\pi}},
\end{aligned} \tag{160}$$

where $k_1 = 1100$ and $k_2 = 3630$. As (160) does not depend on z or $\mathbf{i}$, it bounds $\sup_z P(S \in [z, z + \lambda] | \mathbf{I})$ in (156).

Now using (156), (160) and (150), we obtain

$$\begin{aligned}
A_3 &\leq \frac{1}{\lambda} (2 \max_{l \in \{4, 6, 8\}} \delta^1(k_1\beta_D, n - l) + k_2 \frac{\beta_D}{n} + \frac{2\lambda}{\sqrt{2\pi}}) E|W - W^*| \\
&\leq \frac{1}{\lambda} (2 \max_{l \in \{4, 6, 8\}} \delta^1(k_1\beta_D, n - l) + k_2 \frac{\beta_D}{n} + \frac{2\lambda}{\sqrt{2\pi}}) 113 \frac{\beta_D}{n}.
\end{aligned} \tag{161}$$

Combining (150), (154), (161) and using (149) we obtain for $n \geq 1008$,

$$\begin{aligned}\delta_D &\leq 3842 \frac{\beta_D}{n} + \frac{1}{\lambda} (2 \max_{l \in \{4,6,8\}} \delta^1(k_1 \beta_D, n-l) + k_2 \frac{\beta_D}{n} + \frac{2\lambda}{\sqrt{2\pi}}) 113 \frac{\beta_D}{n} + \frac{\lambda}{\sqrt{2\pi}} \\ &\leq 3842 \frac{\gamma}{n} + \frac{113\gamma}{n\lambda} (2 \max_{l \in \{4,6,8\}} \delta^1(k_1 \beta_D, n-l) + k_2 \frac{\gamma}{n} + \frac{2\lambda}{\sqrt{2\pi}}) + \frac{\lambda}{\sqrt{2\pi}},\end{aligned}$$

since $\beta_D \leq \gamma$ for all $D \in M_n^1(\gamma)$. Setting $\lambda = (113 \times 8)k_1\gamma/n$, we obtain for $n \geq 1008$,

$$\delta_D \leq c \frac{\gamma}{n} + \frac{1}{4} \max_{l \in \{4,6,8\}} \frac{\delta^1(k_1 \beta_D, n-l)}{k_1}, \quad (162)$$

where $c = 400,665$. Since $c > 51/\epsilon_0$, it's clear that if we consider $D \in M_n^1(\gamma)$, with $\beta_D/n > \epsilon_0/51$, then $\delta_D < c\gamma/n$. Hence (162) holds for all $D \in M_n^1(\gamma)$ with $n \geq 1008$. Taking supremum over $D \in M_n^1(\gamma)$, we have, for $n \geq 1008$,

$$\delta^1(\gamma, n) \leq c \frac{\gamma}{n} + \frac{1}{4} \max_{l \in \{4,6,8\}} \frac{\delta^1(k_1 \gamma, n-l)}{k_1},$$

where $c = 400,665$. Now multiplying by n/γ and taking supremum over γ we obtain

$$\sup_{\gamma} n \frac{\delta^1(\gamma, n)}{\gamma} \leq c + \frac{2}{7} \max_{l \in \{4,6,8\}} \sup_{\gamma} (n-l) \frac{\delta^1(\gamma, n-l)}{\gamma} \quad \text{for all } n \geq 1008. \quad (163)$$

If $D \in M_n^1(\gamma)$ and $n \geq 1000$ then (116) shows that $\beta_D \geq 2$, and hence $D \notin M_n^1(\gamma)$ for all $\gamma < 2$. Since $\delta^1(\gamma, n) \leq 1$ for all $n \in \mathbb{N}$, for $1000 \leq n \leq 1008$ we have

$$\sup_{\gamma} n \frac{\delta^1(\gamma, n)}{\gamma} = \sup_{\gamma \geq 2} n \frac{\delta^1(\gamma, n)}{\gamma} \leq 1008 \sup_{\gamma \geq 2} \frac{\delta^1(\gamma, n)}{\gamma} \leq 504.$$

Since $c > 504$, we conclude (163) holds for $n \geq 1000$, that is

$$\sup_{\gamma} n \frac{\delta^1(\gamma, n)}{\gamma} \leq c + \frac{2}{7} \max_{l \in \{4,6,8\}} \sup_{\gamma} (n-l) \frac{\delta^1(\gamma, n-l)}{\gamma} \quad \text{for all } n \geq 1000. \quad (164)$$

Letting $s_n = \sup_{\gamma} n \delta^1(\gamma, n)/\gamma$ and $a_n = s_{n+999}$ for $n \geq 1$, and $a_n = 0$ for $n \leq 0$, (164) gives

$$a_n \leq c + \frac{2}{7} \max_{l \in \{4,6,8\}} a_{n-l} \quad \text{for } n \geq 1.$$

Using Lemma 5.5 with $\alpha = 2/7$, $c = 400,665$ and noting that $a_1 = n \delta^1(\gamma, 1000)/\gamma \leq 500$ since $\gamma \geq 2$, we obtain

$$a_1(1 - 3\alpha) \leq 500(1 - 6/7) < c \quad \text{which yields } b = \max\{c, a_1(1 - 3\alpha)\} = c,$$

and therefore

$$a_n \leq \frac{b}{1 - 3\alpha} = \frac{c}{1 - 3\alpha} = 2804655.$$

This completes the proof of the lemma. □

Lemmas 5.6, 5.3 and remarks following (116) yield

$$\|F_W - \Phi\|_{\infty} \leq K \frac{\beta_D}{n} \quad \text{with } n \geq 1000,$$

where we can take $K = 22 \times 2804655 + 36 = 61,702,446$. This proves Theorem 5.1 and hence completes the proof of Theorem 2.1 as well.

Acknowledgements

The author would like to thank his advisor Prof. Larry Goldstein for suggesting the problem, pointing out several mistakes in earlier working versions of this paper and giving valuable insight.

References

- [1] BOLTHAUSEN, E.(1984). Estimate of the remainder in a combinatorial central limit theorem, *Z. Wahrsch. Verw. Gebiet*, **66**, 379-386.
- [2] CHEN, L.H.Y and SHAO, Q.M.(2005). Stein's method for normal approximation, *An Introduction to Stein's Method*, Chen,L.H.Y and Barbour,A.D. eds,Lecture Notes Series No. 4, Institute for Mathematical Sciences, National University of Singapore, Singapore University Press and World Scientific 2005, 1-59.
- [3] GOLDSTEIN, L. (2007). L^1 bounds in normal approximation, *Ann. Probab.*, **35**, 1888-1930.
- [4] GOLDSTEIN, L. (2005).Berry Esseen Bounds for Combinatorial Central Limit Theorems and Pattern Occurrences, using Zero and Size Biasing in *J. Appl. Probab.*, **42**, 661-683.
- [5] GOLDSTEIN, L. and REINERT, G. (1997).Stein's Method and the Zero Bias Transformation with Application to Simple Random Sampling,*Ann. Appl. Probab.*, **7**, 935-952
- [6] GOLDSTEIN, L. and RINOTT, Y. (2004).A permutation test for matching and its asymptotic distribution, *Metron*, **61**, 375-388.
- [7] HO, S.T. and CHEN, LOUIS H.Y. (1978).An L_p bound for the remainder in a combinatorial central limit theorem, *Ann. Probab.*, **6**, 231-249.
- [8] Hoeffding, W. (1951). A combinatorial central limit theorem, *Ann. Math. Statistics*, **22**, 558-566.
- [9] KOLCHIN, V.F. and CHISTYAKOV, V.P. (1973). On a combinatorial central limit theorem, *Theory Probab. Appl.*, **18**, 728-739.
- [10] MOTOO, M. (1957). On the Hoeffding's combinatorial central limit theorem, *Ann. Inst. Statist. Math.*, **8**, 145-154.
- [11] SCHIFFMAN, A. ET AL (1978). Initial diagnostic hypotheses: factors which may distort physicians judgement, *Organisational Behaviour and Human Performance*, **21**, 305-315.
- [12] STEIN, C. (1972). A bound for the error in the normal approximation to the distribution of a sum of dependent random variables, *Proc. Sixth Berkeley Symp. Math. Statist. Probab.* **2**, 583-602, Univ. California Press, Berkeley.
- [13] VON BAHR, B. (1976).Remainder term estimate in a combinatorial limit theorem., *Z. Wahrsch. Verw. Gebiete*, **35**, 131-139.
- [14] WALD, A. and WOLFOWITZ,J. (1944). Statistical tests based on the permutations of the observations.,*Ann. Math. Statistics*, **15**, 358-372.